

Flow Rates

www.mymathscloud.com

Questions in past papers often come up combined with other topics.
Topic tags have been given for each question to enable you to know if you can do the question or whether you need to wait to cover the additional topic(s).

Scan the QR code(s) or click the link for instant detailed model solutions!

Question 1

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Differentiation, Applications of Integration, Integration

Subtopics: Mean Value Theorem, Interpreting Meaning in Applied Contexts, Rates of Change (Average), Riemann Sums – Trapezoidal Rule, Average Value of a Function, Differentiation Technique – Exponentials, Differentiation Technique – Product Rule

Paper: Part A-Calc / Series: 2001 / Difficulty: Hard / Question Number: 2

t (days)	$W(t)$ (°C)
0	20
3	31
6	28
9	24
12	22
15	21

2. The temperature, in degrees Celsius (°C), of the water in a pond is a differentiable function W of time t . The table above shows the water temperature as recorded every 3 days over a 15-day period.
- (a) Use data from the table to find an approximation for $W'(12)$. Show the computations that lead to your answer. Indicate units of measure.
 - (b) Approximate the average temperature, in degrees Celsius, of the water over the time interval $0 \leq t \leq 15$ days by using a trapezoidal approximation with subintervals of length $\Delta t = 3$ days.
 - (c) A student proposes the function P , given by $P(t) = 20 + 10te^{(-t/3)}$, as a model for the temperature of the water in the pond at time t , where t is measured in days and $P(t)$ is measured in degrees Celsius. Find $P'(12)$. Using appropriate units, explain the meaning of your answer in terms of water temperature.
 - (d) Use the function P defined in part (c) to find the average value, in degrees Celsius, of $P(t)$ over the time interval $0 \leq t \leq 15$ days.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)


Question 2

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Integration

Subtopics: Rates of Change (Average), Riemann Sums – Left, Interpreting Meaning in Applied Contexts, Accumulation of Change

Paper: Part A-Calc / Series: 2003 / Difficulty: Hard / Question Number: 3



t (minutes)	$R(t)$ (gallons per minute)
0	20
30	30
40	40
50	55
70	65
90	70

3. The rate of fuel consumption, in gallons per minute, recorded during an airplane flight is given by a twice-differentiable and strictly increasing function R of time t . The graph of R and a table of selected values of $R(t)$, for the time interval $0 \leq t \leq 90$ minutes, are shown above.

- (a) Use data from the table to find an approximation for $R'(45)$. Show the computations that lead to your answer. Indicate units of measure.
- (b) The rate of fuel consumption is increasing fastest at time $t = 45$ minutes. What is the value of $R''(45)$? Explain your reasoning.
- (c) Approximate the value of $\int_0^{90} R(t) dt$ using a left Riemann sum with the five subintervals indicated by the data in the table. Is this numerical approximation less than the value of $\int_0^{90} R(t) dt$? Explain your reasoning.
- (d) For $0 < b \leq 90$ minutes, explain the meaning of $\int_0^b R(t) dt$ in terms of fuel consumption for the plane. Explain the meaning of $\frac{1}{b} \int_0^b R(t) dt$ in terms of fuel consumption for the plane. Indicate units of measure in both answers.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 3

Qualification: AP Calculus AB

Areas: Applications of Integration, Applications of Differentiation

Subtopics: Total Amount, Increasing/Decreasing , Average Value of a Function, Rates of Change (Average), Accumulation of Change

Paper: Part A-Calc / Series: 2004 / Difficulty: Easy / Question Number: 1

1. Traffic flow is defined as the rate at which cars pass through an intersection, measured in cars per minute. The traffic flow at a particular intersection is modeled by the function F defined by

$$F(t) = 82 + 4 \sin\left(\frac{t}{2}\right) \text{ for } 0 \leq t \leq 30,$$

where $F(t)$ is measured in cars per minute and t is measured in minutes.

- (a) To the nearest whole number, how many cars pass through the intersection over the 30-minute period?
- (b) Is the traffic flow increasing or decreasing at $t = 7$? Give a reason for your answer.
- (c) What is the average value of the traffic flow over the time interval $10 \leq t \leq 15$? Indicate units of measure.
- (d) What is the average rate of change of the traffic flow over the time interval $10 \leq t \leq 15$? Indicate units of measure.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 4

Qualification: AP Calculus AB

Areas: Differentiation, Integration, Applications of Integration, Applications of Differentiation

Subtopics: Rates of Change (Average), Riemann Sums – Trapezoidal Rule, Average Value of a Function, Fundamental Theorem of Calculus (First), Interpreting Meaning in Applied Contexts, Mean Value Theorem

Paper: Part A-Calc / Series: 2005 / Difficulty: Very Hard / Question Number: 3

Distance x (cm)	0	1	5	6	8
Temperature $T(x)$ ($^{\circ}\text{C}$)	100	93	70	62	55

3. A metal wire of length 8 centimeters (cm) is heated at one end. The table above gives selected values of the temperature $T(x)$, in degrees Celsius ($^{\circ}\text{C}$), of the wire x cm from the heated end. The function T is decreasing and twice differentiable.
- (a) Estimate $T'(7)$. Show the work that leads to your answer. Indicate units of measure.
- (b) Write an integral expression in terms of $T(x)$ for the average temperature of the wire. Estimate the average temperature of the wire using a trapezoidal sum with the four subintervals indicated by the data in the table. Indicate units of measure.
- (c) Find $\int_0^8 T'(x) dx$, and indicate units of measure. Explain the meaning of $\int_0^8 T'(x) dx$ in terms of the temperature of the wire.
- (d) Are the data in the table consistent with the assertion that $T''(x) > 0$ for every x in the interval $0 < x < 8$? Explain your answer.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 5

Qualification: AP Calculus AB

Areas: Applications of Integration

Subtopics: Kinematics (Displacement, Velocity, and Acceleration), Rates of Change (Average), Interpreting Meaning in Applied Contexts, Riemann Sums – Midpoint

Paper: Part B-Non-Calc / Series: 2006 / Difficulty: Medium / Question Number: 4

t (seconds)	0	10	20	30	40	50	60	70	80
$v(t)$ (feet per second)	5	14	22	29	35	40	44	47	49

4. Rocket A has positive velocity $v(t)$ after being launched upward from an initial height of 0 feet at time $t = 0$ seconds. The velocity of the rocket is recorded for selected values of t over the interval $0 \leq t \leq 80$ seconds, as shown in the table above.
- (a) Find the average acceleration of rocket A over the time interval $0 \leq t \leq 80$ seconds. Indicate units of measure.
- (b) Using correct units, explain the meaning of $\int_{10}^{70} v(t) dt$ in terms of the rocket's flight. Use a midpoint Riemann sum with 3 subintervals of equal length to approximate $\int_{10}^{70} v(t) dt$.
- (c) Rocket B is launched upward with an acceleration of $a(t) = \frac{3}{\sqrt{t+1}}$ feet per second per second. At time $t = 0$ seconds, the initial height of the rocket is 0 feet, and the initial velocity is 2 feet per second. Which of the two rockets is traveling faster at time $t = 80$ seconds? Explain your answer.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 6

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Integration

Subtopics: Tangents To Curves, Concavity, Rates of Change (Instantaneous), Riemann Sums – Right, Interpreting Meaning in Applied Contexts

Paper: Part B-Non-Calc / Series: 2007 / Difficulty: Hard / Question Number: 5

t (minutes)	0	2	5	7	11	12
$r'(t)$ (feet per minute)	5.7	4.0	2.0	1.2	0.6	0.5

5. The volume of a spherical hot air balloon expands as the air inside the balloon is heated. The radius of the balloon, in feet, is modeled by a twice-differentiable function r of time t , where t is measured in minutes. For $0 < t < 12$, the graph of r is concave down. The table above gives selected values of the rate of change, $r'(t)$, of the radius of the balloon over the time interval $0 \leq t \leq 12$. The radius of the balloon is 30 feet when $t = 5$.

(Note: The volume of a sphere of radius r is given by $V = \frac{4}{3}\pi r^3$.)

- (a) Estimate the radius of the balloon when $t = 5.4$ using the tangent line approximation at $t = 5$. Is your estimate greater than or less than the true value? Give a reason for your answer.
- (b) Find the rate of change of the volume of the balloon with respect to time when $t = 5$. Indicate units of measure.
- (c) Use a right Riemann sum with the five subintervals indicated by the data in the table to approximate $\int_0^{12} r'(t) dt$. Using correct units, explain the meaning of $\int_0^{12} r'(t) dt$ in terms of the radius of the balloon.
- (d) Is your approximation in part (c) greater than or less than $\int_0^{12} r'(t) dt$? Give a reason for your answer.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 7

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Differentiation

Subtopics: Rates of Change (Instantaneous), Rates of Change (Average), Differentiation Technique – Chain Rule, Interpreting Meaning in Applied Contexts, Modelling Situations

Paper: Part A-Calc / Series: 2007-Form-B / Difficulty: Medium / Question Number: 3

3. The wind chill is the temperature, in degrees Fahrenheit ($^{\circ}\text{F}$), a human feels based on the air temperature, in degrees Fahrenheit, and the wind velocity v , in miles per hour (mph). If the air temperature is 32°F , then the wind chill is given by $W(v) = 55.6 - 22.1v^{0.16}$ and is valid for $5 \leq v \leq 60$.
- (a) Find $W'(20)$. Using correct units, explain the meaning of $W'(20)$ in terms of the wind chill.
- (b) Find the average rate of change of W over the interval $5 \leq v \leq 60$. Find the value of v at which the instantaneous rate of change of W is equal to the average rate of change of W over the interval $5 \leq v \leq 60$.
- (c) Over the time interval $0 \leq t \leq 4$ hours, the air temperature is a constant 32°F . At time $t = 0$, the wind velocity is $v = 20$ mph. If the wind velocity increases at a constant rate of 5 mph per hour, what is the rate of change of the wind chill with respect to time at $t = 3$ hours? Indicate units of measure.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 8

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Applications of Integration

Subtopics: Rates of Change (Average), Riemann Sums – Trapezoidal Rule, Mean Value Theorem, Intermediate Value Theorem, Local or Relative Minima and Maxima, Total Amount

Paper: Part A-Calc / Series: 2008 / Difficulty: Hard / Question Number: 2

t (hours)	0	1	3	4	7	8	9
$L(t)$ (people)	120	156	176	126	150	80	0

2. Concert tickets went on sale at noon ($t = 0$) and were sold out within 9 hours. The number of people waiting in line to purchase tickets at time t is modeled by a twice-differentiable function L for $0 \leq t \leq 9$. Values of $L(t)$ at various times t are shown in the table above.
- (a) Use the data in the table to estimate the rate at which the number of people waiting in line was changing at 5:30 P.M. ($t = 5.5$). Show the computations that lead to your answer. Indicate units of measure.
- (b) Use a trapezoidal sum with three subintervals to estimate the average number of people waiting in line during the first 4 hours that tickets were on sale.
- (c) For $0 \leq t \leq 9$, what is the fewest number of times at which $L'(t)$ must equal 0? Give a reason for your answer.
- (d) The rate at which tickets were sold for $0 \leq t \leq 9$ is modeled by $r(t) = 550te^{-t/2}$ tickets per hour. Based on the model, how many tickets were sold by 3 P.M. ($t = 3$), to the nearest whole number?
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 9

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Differentiation

Subtopics: Rates of Change (Instantaneous), Implicit Differentiation, Modelling Situations, Global or Absolute Minima and Maxima, Related Rates

Paper: Part A-Calc / Series: 2008 / Difficulty: Medium / Question Number: 3

3. Oil is leaking from a pipeline on the surface of a lake and forms an oil slick whose volume increases at a constant rate of 2000 cubic centimeters per minute. The oil slick takes the form of a right circular cylinder with both its radius and height changing with time. (Note: The volume V of a right circular cylinder with radius r and height h is given by $V = \pi r^2 h$.)
- (a) At the instant when the radius of the oil slick is 100 centimeters and the height is 0.5 centimeter, the radius is increasing at the rate of 2.5 centimeters per minute. At this instant, what is the rate of change of the height of the oil slick with respect to time, in centimeters per minute?
- (b) A recovery device arrives on the scene and begins removing oil. The rate at which oil is removed is $R(t) = 400\sqrt{t}$ cubic centimeters per minute, where t is the time in minutes since the device began working. Oil continues to leak at the rate of 2000 cubic centimeters per minute. Find the time t when the oil slick reaches its maximum volume. Justify your answer.
- (c) By the time the recovery device began removing oil, 60,000 cubic centimeters of oil had already leaked. Write, but do not evaluate, an expression involving an integral that gives the volume of oil at the time found in part (b).
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 10

Qualification: AP Calculus AB

Areas: Applications of Integration, Differentiation

Subtopics: Kinematics (Displacement, Velocity, and Acceleration), Rates of Change (Instantaneous), Differentiation Technique – Chain Rule, Total Amount

Paper: Part A-Calc / Series: 2008-Form-B / Difficulty: Medium / Question Number: 2

2. For time $t \geq 0$ hours, let $r(t) = 120(1 - e^{-10t^2})$ represent the speed, in kilometers per hour, at which a car travels along a straight road. The number of liters of gasoline used by the car to travel x kilometers is modeled by $g(x) = 0.05x(1 - e^{-x/2})$.
- (a) How many kilometers does the car travel during the first 2 hours?
 - (b) Find the rate of change with respect to time of the number of liters of gasoline used by the car when $t = 2$ hours. Indicate units of measure.
 - (c) How many liters of gasoline have been used by the car when it reaches a speed of 80 kilometers per hour?

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 11

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Integration

Subtopics: Rates of Change (Average), Fundamental Theorem of Calculus (First), Riemann Sums – Left

Paper: Part B-Non-Calc / Series: 2009 / Difficulty: Medium / Question Number: 5

x	2	3	5	8	13
$f(x)$	1	4	-2	3	6

5. Let f be a function that is twice differentiable for all real numbers. The table above gives values of f for selected points in the closed interval $2 \leq x \leq 13$.

(a) Estimate $f'(4)$. Show the work that leads to your answer.

(b) Evaluate $\int_2^{13} (3 - 5f'(x)) dx$. Show the work that leads to your answer.

(c) Use a left Riemann sum with subintervals indicated by the data in the table to approximate $\int_2^{13} f(x) dx$.
Show the work that leads to your answer.

(d) Suppose $f'(5) = 3$ and $f''(x) < 0$ for all x in the closed interval $5 \leq x \leq 8$. Use the line tangent to the graph of f at $x = 5$ to show that $f(7) \leq 4$. Use the secant line for the graph of f on $5 \leq x \leq 8$ to show that $f(7) \geq \frac{4}{3}$.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 12

Qualification: AP Calculus AB

Areas: Integration

Subtopics: Kinematics (Displacement, Velocity, and Acceleration), Interpreting Meaning in Applied Contexts, Riemann Sums – Trapezoidal Rule, Derivative Tables, Rates of Change (Average)

Paper: Part B-Non-Calc / Series: 2009-Form-B / Difficulty: Somewhat Challenging / Question Number: 6

t (seconds)	0	8	20	25	32	40
$v(t)$ (meters per second)	3	5	-10	-8	-4	7

6. The velocity of a particle moving along the x -axis is modeled by a differentiable function v , where the position x is measured in meters, and time t is measured in seconds. Selected values of $v(t)$ are given in the table above. The particle is at position $x = 7$ meters when $t = 0$ seconds.
- (a) Estimate the acceleration of the particle at $t = 36$ seconds. Show the computations that lead to your answer. Indicate units of measure.
- (b) Using correct units, explain the meaning of $\int_{20}^{40} v(t) dt$ in the context of this problem. Use a trapezoidal sum with the three subintervals indicated by the data in the table to approximate $\int_{20}^{40} v(t) dt$.
- (c) For $0 \leq t \leq 40$, must the particle change direction in any of the subintervals indicated by the data in the table? If so, identify the subintervals and explain your reasoning. If not, explain why not.
- (d) Suppose that the acceleration of the particle is positive for $0 < t < 8$ seconds. Explain why the position of the particle at $t = 8$ seconds must be greater than $x = 30$ meters.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 13

Qualification: AP Calculus AB

Areas: Applications of Integration, Applications of Differentiation

Subtopics: Rates of Change (Average), Riemann Sums – Trapezoidal Rule, Interpreting Meaning in Applied Contexts, Modelling Situations, Global or Absolute Minima and Maxima

Paper: Part A-Calc / Series: 2010 / Difficulty: Medium / Question Number: 2

t (hours)	0	2	5	7	8
$E(t)$ (hundreds of entries)	0	4	13	21	23

2. A zoo sponsored a one-day contest to name a new baby elephant. Zoo visitors deposited entries in a special box between noon ($t = 0$) and 8 P.M. ($t = 8$). The number of entries in the box t hours after noon is modeled by a differentiable function E for $0 \leq t \leq 8$. Values of $E(t)$, in hundreds of entries, at various times t are shown in the table above.

- (a) Use the data in the table to approximate the rate, in hundreds of entries per hour, at which entries were being deposited at time $t = 6$. Show the computations that lead to your answer.

- (b) Use a trapezoidal sum with the four subintervals given by the table to approximate the value of $\frac{1}{8} \int_0^8 E(t) dt$.

Using correct units, explain the meaning of $\frac{1}{8} \int_0^8 E(t) dt$ in terms of the number of entries.

- (c) At 8 P.M., volunteers began to process the entries. They processed the entries at a rate modeled by the function P , where $P(t) = t^3 - 30t^2 + 298t - 976$ hundreds of entries per hour for $8 \leq t \leq 12$. According to the model, how many entries had not yet been processed by midnight ($t = 12$)?
- (d) According to the model from part (c), at what time were the entries being processed most quickly? Justify your answer.
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 14

Qualification: AP Calculus AB

Areas: Applications of Integration, Integration

Subtopics: Rates of Change (Average), Riemann Sums – Trapezoidal Rule, Interpreting Meaning in Applied Contexts, Fundamental Theorem of Calculus (First)

Paper: Part A-Calc / Series: 2011 / Difficulty: Easy / Question Number: 2

t (minutes)	0	2	5	9	10
$H(t)$ (degrees Celsius)	66	60	52	44	43

2. As a pot of tea cools, the temperature of the tea is modeled by a differentiable function H for $0 \leq t \leq 10$, where time t is measured in minutes and temperature $H(t)$ is measured in degrees Celsius. Values of $H(t)$ at selected values of time t are shown in the table above.
- (a) Use the data in the table to approximate the rate at which the temperature of the tea is changing at time $t = 3.5$. Show the computations that lead to your answer.
- (b) Using correct units, explain the meaning of $\frac{1}{10} \int_0^{10} H(t) dt$ in the context of this problem. Use a trapezoidal sum with the four subintervals indicated by the table to estimate $\frac{1}{10} \int_0^{10} H(t) dt$.
- (c) Evaluate $\int_0^{10} H'(t) dt$. Using correct units, explain the meaning of the expression in the context of this problem.
- (d) At time $t = 0$, biscuits with temperature 100°C were removed from an oven. The temperature of the biscuits at time t is modeled by a differentiable function B for which it is known that $B'(t) = -13.84e^{-0.173t}$. Using the given models, at time $t = 10$, how much cooler are the biscuits than the tea?
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 15

Qualification: AP Calculus AB

Areas: Applications of Differentiation

Subtopics: Rates of Change (Average), Modelling Situations, Intermediate Value Theorem

Paper: Part A-Calc / Series: 2011-Form-B / Difficulty: Easy / Question Number: 1

1. A cylindrical can of radius 10 millimeters is used to measure rainfall in Stormville. The can is initially empty, and rain enters the can during a 60-day period. The height of water in the can is modeled by the function S , where $S(t)$ is measured in millimeters and t is measured in days for $0 \leq t \leq 60$. The rate at which the height of the water is rising in the can is given by $S'(t) = 2 \sin(0.03t) + 1.5$.
- (a) According to the model, what is the height of the water in the can at the end of the 60-day period?
- (b) According to the model, what is the average rate of change in the height of water in the can over the 60-day period? Show the computations that lead to your answer. Indicate units of measure.
- (c) Assuming no evaporation occurs, at what rate is the volume of water in the can changing at time $t = 7$? Indicate units of measure.
- (d) During the same 60-day period, rain on Monsoon Mountain accumulates in a can identical to the one in Stormville. The height of the water in the can on Monsoon Mountain is modeled by the function M , where $M(t) = \frac{1}{400}(3t^3 - 30t^2 + 330t)$. The height $M(t)$ is measured in millimeters, and t is measured in days for $0 \leq t \leq 60$. Let $D(t) = M'(t) - S'(t)$. Apply the Intermediate Value Theorem to the function D on the interval $0 \leq t \leq 60$ to justify that there exists a time t , $0 < t < 60$, at which the heights of water in the two cans are changing at the same rate.
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 16

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Differentiation

Subtopics: Kinematics (Displacement, Velocity, and Acceleration), Rates of Change (Average), Riemann Sums – Left, Mean Value Theorem, Implicit Differentiation

Paper: Part B-Non-Calc / Series: 2011-Form-B / Difficulty: Medium / Question Number: 5

t (seconds)	0	10	40	60
$B(t)$ (meters)	100	136	9	49
$v(t)$ (meters per second)	2.0	2.3	2.5	4.6

5. Ben rides a unicycle back and forth along a straight east-west track. The twice-differentiable function B models Ben's position on the track, measured in meters from the western end of the track, at time t , measured in seconds from the start of the ride. The table above gives values for $B(t)$ and Ben's velocity, $v(t)$, measured in meters per second, at selected times t .
- (a) Use the data in the table to approximate Ben's acceleration at time $t = 5$ seconds. Indicate units of measure.
- (b) Using correct units, interpret the meaning of $\int_0^{60} |v(t)| dt$ in the context of this problem. Approximate $\int_0^{60} |v(t)| dt$ using a left Riemann sum with the subintervals indicated by the data in the table.
- (c) For $40 \leq t \leq 60$, must there be a time t when Ben's velocity is 2 meters per second? Justify your answer.
- (d) A light is directly above the western end of the track. Ben rides so that at time t , the distance $L(t)$ between Ben and the light satisfies $(L(t))^2 = 12^2 + (B(t))^2$. At what rate is the distance between Ben and the light changing at time $t = 40$?
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 17

Qualification: AP Calculus AB

Areas: Applications of Integration, Integration, Differentiation

Subtopics: Rates of Change (Average), Interpreting Meaning in Applied Contexts, Fundamental Theorem of Calculus (First), Riemann Sums – Left

Paper: Part A-Calc / Series: 2012 / Difficulty: Medium / Question Number: 1

t (minutes)	0	4	9	15	20
$W(t)$ (degrees Fahrenheit)	55.0	57.1	61.8	67.9	71.0

1. The temperature of water in a tub at time t is modeled by a strictly increasing, twice-differentiable function W , where $W(t)$ is measured in degrees Fahrenheit and t is measured in minutes. At time $t = 0$, the temperature of the water is 55°F . The water is heated for 30 minutes, beginning at time $t = 0$. Values of $W(t)$ at selected times t for the first 20 minutes are given in the table above.
- (a) Use the data in the table to estimate $W'(12)$. Show the computations that lead to your answer. Using correct units, interpret the meaning of your answer in the context of this problem.
- (b) Use the data in the table to evaluate $\int_0^{20} W'(t) dt$. Using correct units, interpret the meaning of $\int_0^{20} W'(t) dt$ in the context of this problem.
- (c) For $0 \leq t \leq 20$, the average temperature of the water in the tub is $\frac{1}{20} \int_0^{20} W(t) dt$. Use a left Riemann sum with the four subintervals indicated by the data in the table to approximate $\frac{1}{20} \int_0^{20} W(t) dt$. Does this approximation overestimate or underestimate the average temperature of the water over these 20 minutes? Explain your reasoning.
- (d) For $20 \leq t \leq 25$, the function W that models the water temperature has first derivative given by $W'(t) = 0.4\sqrt{t} \cos(0.06t)$. Based on the model, what is the temperature of the water at time $t = 25$?
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 18

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Integration

Subtopics: Rates of Change (Average), Mean Value Theorem, Riemann Sums – Midpoint, Interpreting Meaning in Applied Contexts, Rates of Change (Instantaneous)

Paper: Part B-Non-Calc / Series: 2013 / Difficulty: Medium / Question Number: 3

t (minutes)	0	1	2	3	4	5	6
$C(t)$ (ounces)	0	5.3	8.8	11.2	12.8	13.8	14.5

3. Hot water is dripping through a coffeemaker, filling a large cup with coffee. The amount of coffee in the cup at time t , $0 \leq t \leq 6$, is given by a differentiable function C , where t is measured in minutes. Selected values of $C(t)$, measured in ounces, are given in the table above.
- (a) Use the data in the table to approximate $C'(3.5)$. Show the computations that lead to your answer, and indicate units of measure.
- (b) Is there a time t , $2 \leq t \leq 4$, at which $C'(t) = 2$? Justify your answer.
- (c) Use a midpoint sum with three subintervals of equal length indicated by the data in the table to approximate the value of $\frac{1}{6} \int_0^6 C(t) dt$. Using correct units, explain the meaning of $\frac{1}{6} \int_0^6 C(t) dt$ in the context of the problem.
- (d) The amount of coffee in the cup, in ounces, is modeled by $B(t) = 16 - 16e^{-0.4t}$. Using this model, find the rate at which the amount of coffee in the cup is changing when $t = 5$.
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 19

Qualification: AP Calculus AB

Areas: Applications of Integration, Differentiation

Subtopics: Rates of Change (Average), Interpreting Meaning in Applied Contexts, Total Amount, Average Value of a Function, Modelling Situations

Paper: Part A-Calc / Series: 2014 / Difficulty: Medium / Question Number: 1

1. Grass clippings are placed in a bin, where they decompose. For $0 \leq t \leq 30$, the amount of grass clippings remaining in the bin is modeled by $A(t) = 6.687(0.931)^t$, where $A(t)$ is measured in pounds and t is measured in days.
- (a) Find the average rate of change of $A(t)$ over the interval $0 \leq t \leq 30$. Indicate units of measure.
 - (b) Find the value of $A'(15)$. Using correct units, interpret the meaning of the value in the context of the problem.
 - (c) Find the time t for which the amount of grass clippings in the bin is equal to the average amount of grass clippings in the bin over the interval $0 \leq t \leq 30$.
 - (d) For $t > 30$, $L(t)$, the linear approximation to A at $t = 30$, is a better model for the amount of grass clippings remaining in the bin. Use $L(t)$ to predict the time at which there will be 0.5 pound of grass clippings remaining in the bin. Show the work that leads to your answer.
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 20

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Integration

Subtopics: Rates of Change (Average), Intermediate Value Theorem, Riemann Sums – Trapezoidal Rule, Rates of Change (Instantaneous), Modelling Situations, Implicit Differentiation

Paper: Part B-Non-Calc / Series: 2014 / Difficulty: Somewhat Challenging / Question Number: 4

t (minutes)	0	2	5	8	12
$v_A(t)$ (meters/minute)	0	100	40	-120	-150

4. Train A runs back and forth on an east-west section of railroad track. Train A 's velocity, measured in meters per minute, is given by a differentiable function $v_A(t)$, where time t is measured in minutes. Selected values for $v_A(t)$ are given in the table above.
- (a) Find the average acceleration of train A over the interval $2 \leq t \leq 8$.
- (b) Do the data in the table support the conclusion that train A 's velocity is -100 meters per minute at some time t with $5 < t < 8$? Give a reason for your answer.
- (c) At time $t = 2$, train A 's position is 300 meters east of the Origin Station, and the train is moving to the east. Write an expression involving an integral that gives the position of train A , in meters from the Origin Station, at time $t = 12$. Use a trapezoidal sum with three subintervals indicated by the table to approximate the position of the train at time $t = 12$.
- (d) A second train, train B , travels north from the Origin Station. At time t the velocity of train B is given by $v_B(t) = -5t^2 + 60t + 25$, and at time $t = 2$ the train is 400 meters north of the station. Find the rate, in meters per minute, at which the distance between train A and train B is changing at time $t = 2$.
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 21

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Applications of Integration

Subtopics: Rates of Change (Average), Interpreting Meaning in Applied Contexts, Riemann Sums – Right, Kinematics (Displacement, Velocity, and Acceleration), Average Value of a Function

Paper: Part B-Non-Calc / Series: 2015 / Difficulty: Medium / Question Number: 3

t (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	-220	150

3. Johanna jogs along a straight path. For $0 \leq t \leq 40$, Johanna's velocity is given by a differentiable function v . Selected values of $v(t)$, where t is measured in minutes and $v(t)$ is measured in meters per minute, are given in the table above.

(a) Use the data in the table to estimate the value of $v'(16)$.

(b) Using correct units, explain the meaning of the definite integral $\int_0^{40} |v(t)| dt$ in the context of the problem.

Approximate the value of $\int_0^{40} |v(t)| dt$ using a right Riemann sum with the four subintervals indicated in the table.

(c) Bob is riding his bicycle along the same path. For $0 \leq t \leq 10$, Bob's velocity is modeled by

$$B(t) = t^3 - 6t^2 + 300, \text{ where } t \text{ is measured in minutes and } B(t) \text{ is measured in meters per minute.}$$

Find Bob's acceleration at time $t = 5$.

(d) Based on the model B from part (c), find Bob's average velocity during the interval $0 \leq t \leq 10$.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 22

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Integration

Subtopics: Rates of Change (Average), Riemann Sums – Left , Increasing/Decreasing , Total Amount, Intermediate Value Theorem, Modelling Situations

Paper: Part A-Calc / Series: 2016 / Difficulty: Somewhat Challenging / Question Number: 1

t (hours)	0	1	3	6	8
$R(t)$ (liters / hour)	1340	1190	950	740	700

1. Water is pumped into a tank at a rate modeled by $W(t) = 2000e^{-t^2/20}$ liters per hour for $0 \leq t \leq 8$, where t is measured in hours. Water is removed from the tank at a rate modeled by $R(t)$ liters per hour, where R is differentiable and decreasing on $0 \leq t \leq 8$. Selected values of $R(t)$ are shown in the table above. At time $t = 0$, there are 50,000 liters of water in the tank.
- (a) Estimate $R'(2)$. Show the work that leads to your answer. Indicate units of measure.
- (b) Use a left Riemann sum with the four subintervals indicated by the table to estimate the total amount of water removed from the tank during the 8 hours. Is this an overestimate or an underestimate of the total amount of water removed? Give a reason for your answer.
- (c) Use your answer from part (b) to find an estimate of the total amount of water in the tank, to the nearest liter, at the end of 8 hours.
- (d) For $0 \leq t \leq 8$, is there a time t when the rate at which water is pumped into the tank is the same as the rate at which water is removed from the tank? Explain why or why not.
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 23

Qualification: AP Calculus AB

Areas: Applications of Integration, Integration

Subtopics: Riemann Sums – Left , Increasing/Decreasing , Modelling Situations, Differentiation Technique – Chain Rule, Related Rates

Paper: Part A-Calc / Series: 2017 / Difficulty: Medium / Question Number: 1

h (feet)	0	2	5	10
$A(h)$ (square feet)	50.3	14.4	6.5	2.9

1. A tank has a height of 10 feet. The area of the horizontal cross section of the tank at height h feet is given by the function A , where $A(h)$ is measured in square feet. The function A is continuous and decreases as h increases. Selected values for $A(h)$ are given in the table above.
- (a) Use a left Riemann sum with the three subintervals indicated by the data in the table to approximate the volume of the tank. Indicate units of measure.
- (b) Does the approximation in part (a) overestimate or underestimate the volume of the tank? Explain your reasoning.
- (c) The area, in square feet, of the horizontal cross section at height h feet is modeled by the function f given by $f(h) = \frac{50.3}{e^{0.2h} + h}$. Based on this model, find the volume of the tank. Indicate units of measure.
- (d) Water is pumped into the tank. When the height of the water is 5 feet, the height is increasing at the rate of 0.26 foot per minute. Using the model from part (c), find the rate at which the volume of water is changing with respect to time when the height of the water is 5 feet. Indicate units of measure.
-

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 24

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Applications of Integration

Subtopics: Rates of Change (Average), Mean Value Theorem, Average Value of a Function, Riemann Sums – Trapezoidal Rule, Modelling Situations, Rates of Change (Instantaneous), Related Rates

Paper: Part B-Non-Calc / Series: 2018 / Difficulty: Medium / Question Number: 4

t (years)	2	3	5	7	10
$H(t)$ (meters)	1.5	2	6	11	15

4. The height of a tree at time t is given by a twice-differentiable function H , where $H(t)$ is measured in meters and t is measured in years. Selected values of $H(t)$ are given in the table above.
- (a) Use the data in the table to estimate $H'(6)$. Using correct units, interpret the meaning of $H'(6)$ in the context of the problem.
- (b) Explain why there must be at least one time t , for $2 < t < 10$, such that $H'(t) = 2$.
- (c) Use a trapezoidal sum with the four subintervals indicated by the data in the table to approximate the average height of the tree over the time interval $2 \leq t \leq 10$.
- (d) The height of the tree, in meters, can also be modeled by the function G , given by $G(x) = \frac{100x}{1+x}$, where x is the diameter of the base of the tree, in meters. When the tree is 50 meters tall, the diameter of the base of the tree is increasing at a rate of 0.03 meter per year. According to this model, what is the rate of change of the height of the tree with respect to time, in meters per year, at the time when the tree is 50 meters tall?

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 25

Qualification: AP Calculus AB

Areas: Applications of Integration, Applications of Differentiation, Integration

Subtopics: Interpreting Meaning in Applied Contexts, Rates of Change (Average), Riemann Sums – Right, Increasing/Decreasing, Differentiation Technique – Chain Rule, Average Value of a Function

Paper: Part A-Calc / Series: 2021 / Difficulty: Somewhat Challenging / Question Number: 1

r (centimeters)	0	1	2	2.5	4
$f(r)$ (milligrams per square centimeter)	1	2	6	10	18

1. The density of a bacteria population in a circular petri dish at a distance r centimeters from the center of the dish is given by an increasing, differentiable function f , where $f(r)$ is measured in milligrams per square centimeter. Values of $f(r)$ for selected values of r are given in the table above.
- (a) Use the data in the table to estimate $f'(2.25)$. Using correct units, interpret the meaning of your answer in the context of this problem.
- (b) The total mass, in milligrams, of bacteria in the petri dish is given by the integral expression $2\pi \int_0^4 r f(r) dr$. Approximate the value of $2\pi \int_0^4 r f(r) dr$ using a right Riemann sum with the four subintervals indicated by the data in the table.
- (c) Is the approximation found in part (b) an overestimate or underestimate of the total mass of bacteria in the petri dish? Explain your reasoning.
- (d) The density of bacteria in the petri dish, for $1 \leq r \leq 4$, is modeled by the function g defined by $g(r) = 2 - 16(\cos(1.57\sqrt{r}))^3$. For what value of k , $1 < k < 4$, is $g(k)$ equal to the average value of $g(r)$ on the interval $1 \leq r \leq 4$?

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 26

Qualification: AP Calculus AB

Areas: Applications of Differentiation, Differentiation, Integration

Subtopics: Rates of Change (Average), Intermediate Value Theorem, Riemann Sums – Right, Implicit Differentiation, Rates of Change (Instantaneous), Differentiation Technique – Chain Rule, Related Rates

Paper: Part B-Non-Calc / Series: 2022 / Difficulty: Medium / Question Number: 4

t (days)	0	3	7	10	12
$r'(t)$ (centimeters per day)	-6.1	-5.0	-4.4	-3.8	-3.5

4. An ice sculpture melts in such a way that it can be modeled as a cone that maintains a conical shape as it decreases in size. The radius of the base of the cone is given by a twice-differentiable function r , where $r(t)$ is measured in centimeters and t is measured in days. The table above gives selected values of $r'(t)$, the rate of change of the radius, over the time interval $0 \leq t \leq 12$.
- (a) Approximate $r''(8.5)$ using the average rate of change of r' over the interval $7 \leq t \leq 10$. Show the computations that lead to your answer, and indicate units of measure.
- (b) Is there a time t , $0 \leq t \leq 3$, for which $r'(t) = -6$? Justify your answer.
- (c) Use a right Riemann sum with the four subintervals indicated in the table to approximate the value of $\int_0^{12} r'(t) dt$.
- (d) The height of the cone decreases at a rate of 2 centimeters per day. At time $t = 3$ days, the radius is 100 centimeters and the height is 50 centimeters. Find the rate of change of the volume of the cone with respect to time, in cubic centimeters per day, at time $t = 3$ days. (The volume V of a cone with radius r and height h is $V = \frac{1}{3}\pi r^2 h$.)

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)

Question 27

Qualification: AP Calculus AB

Areas: Applications of Integration, Applications of Differentiation

Subtopics: Rates of Change (Average), Riemann Sums – Left, Interpreting Meaning in Applied Contexts, Increasing/Decreasing

Paper: Part A-Calc / Series: 2024 / Difficulty: Easy / Question Number: 1

t (minutes)	0	3	7	12
$C(t)$ (degrees Celsius)	100	85	69	55

1. The temperature of coffee in a cup at time t minutes is modeled by a decreasing differentiable function C , where $C(t)$ is measured in degrees Celsius. For $0 \leq t \leq 12$, selected values of $C(t)$ are given in the table shown.
- (a) Approximate $C'(5)$ using the average rate of change of C over the interval $3 \leq t \leq 7$. Show the work that leads to your answer and include units of measure.
- (b) Use a left Riemann sum with the three subintervals indicated by the data in the table to approximate the value of $\int_0^{12} C(t) dt$. Interpret the meaning of $\frac{1}{12} \int_0^{12} C(t) dt$ in the context of the problem.
- (c) For $12 \leq t \leq 20$, the rate of change of the temperature of the coffee is modeled by $C'(t) = \frac{-24.55e^{0.01t}}{t}$, where $C'(t)$ is measured in degrees Celsius per minute. Find the temperature of the coffee at time $t = 20$. Show the setup for your calculations.
- (d) For the model defined in part (c), it can be shown that $C''(t) = \frac{0.2455e^{0.01t}(100 - t)}{t^2}$. For $12 < t < 20$, determine whether the temperature of the coffee is changing at a decreasing rate or at an increasing rate. Give a reason for your answer.

SCAN ME!



Mark Scheme

[View Online](#)

SCAN ME!



Written Mark Scheme

[View Online](#)