



---

# AP<sup>®</sup> Calculus AB 2014 Scoring Guidelines

---

© 2014 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: [www.collegeboard.org](http://www.collegeboard.org).

AP Central is the official online home for the AP Program: [apcentral.collegeboard.org](http://apcentral.collegeboard.org).

**AP<sup>®</sup> CALCULUS AB/CALCULUS BC  
2014 SCORING GUIDELINES**

**Question 1**

Grass clippings are placed in a bin, where they decompose. For  $0 \leq t \leq 30$ , the amount of grass clippings remaining in the bin is modeled by  $A(t) = 6.687(0.931)^t$ , where  $A(t)$  is measured in pounds and  $t$  is measured in days.

- (a) Find the average rate of change of  $A(t)$  over the interval  $0 \leq t \leq 30$ . Indicate units of measure.
- (b) Find the value of  $A'(15)$ . Using correct units, interpret the meaning of the value in the context of the problem.
- (c) Find the time  $t$  for which the amount of grass clippings in the bin is equal to the average amount of grass clippings in the bin over the interval  $0 \leq t \leq 30$ .
- (d) For  $t > 30$ ,  $L(t)$ , the linear approximation to  $A$  at  $t = 30$ , is a better model for the amount of grass clippings remaining in the bin. Use  $L(t)$  to predict the time at which there will be 0.5 pound of grass clippings remaining in the bin. Show the work that leads to your answer.

(a)  $\frac{A(30) - A(0)}{30 - 0} = -0.197$  (or  $-0.196$ ) lbs/day

1 : answer with units

(b)  $A'(15) = -0.164$  (or  $-0.163$ )

The amount of grass clippings in the bin is decreasing at a rate of 0.164 (or 0.163) lbs/day at time  $t = 15$  days.

2 :  $\begin{cases} 1 : A'(15) \\ 1 : \text{interpretation} \end{cases}$

(c)  $A(t) = \frac{1}{30} \int_0^{30} A(t) dt \Rightarrow t = 12.415$  (or 12.414)

2 :  $\begin{cases} 1 : \frac{1}{30} \int_0^{30} A(t) dt \\ 1 : \text{answer} \end{cases}$

(d)  $L(t) = A(30) + A'(30) \cdot (t - 30)$

$A'(30) = -0.055976$

$A(30) = 0.782928$


$L(t) = 0.5 \Rightarrow t = 35.054$

4 :  $\begin{cases} 2 : \text{expression for } L(t) \\ 1 : L(t) = 0.5 \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB  
2014 SCORING GUIDELINES**

**Question 2**

Let  $R$  be the region enclosed by the graph of  $f(x) = x^4 - 2.3x^3 + 4$  and the horizontal line  $y = 4$ , as shown in the figure above.



- (a) Find the volume of the solid generated when  $R$  is rotated about the horizontal line  $y = -2$ .
- (b) Region  $R$  is the base of a solid. For this solid, each cross section perpendicular to the  $x$ -axis is an isosceles right triangle with a leg in  $R$ . Find the volume of the solid.
- (c) The vertical line  $x = k$  divides  $R$  into two regions with equal areas. Write, but do not solve, an equation involving integral expressions whose solution gives the value  $k$ .

(a)  $f(x) = 4 \Rightarrow x = 0, 2.3$

$$\begin{aligned} \text{Volume} &= \pi \int_0^{2.3} [(4 + 2)^2 - (f(x) + 2)^2] dx \\ &= 98.868 \text{ (or } 98.867) \end{aligned}$$

4 :  $\begin{cases} 2 : \text{integrand} \\ 1 : \text{limits} \\ 1 : \text{answer} \end{cases}$

(b)  $\text{Volume} = \int_0^{2.3} \frac{1}{2} (4 - f(x))^2 dx$   
 $= 3.574 \text{ (or } 3.573)$

3 :  $\begin{cases} 2 : \text{integrand} \\ 1 : \text{answer} \end{cases}$

(c)  $\int_0^k (4 - f(x)) dx = \int_k^{2.3} (4 - f(x)) dx$

2 :  $\begin{cases} 1 : \text{area of one region} \\ 1 : \text{equation} \end{cases}$


**AP<sup>®</sup> CALCULUS AB/CALCULUS BC  
2014 SCORING GUIDELINES**

**Question 3**

The function  $f$  is defined on the closed interval  $[-5, 4]$ . The graph of  $f$  consists of three line segments and is shown in the figure above.

Let  $g$  be the function defined by  $g(x) = \int_{-3}^x f(t) dt$ .

- (a) Find  $g(3)$ .
- (b) On what open intervals contained in  $-5 < x < 4$  is the graph of  $g$  both increasing and concave down? Give a reason for your answer.
- (c) The function  $h$  is defined by  $h(x) = \frac{g(x)}{5x}$ . Find  $h'(3)$ .
- (d) The function  $p$  is defined by  $p(x) = f(x^2 - x)$ . Find the slope of the line tangent to the graph of  $p$  at the point where  $x = -1$ .



Graph of  $f$

(a)  $g(3) = \int_{-3}^3 f(t) dt = 6 + 4 - 1 = 9$

1 : answer

(b)  $g'(x) = f(x)$

The graph of  $g$  is increasing and concave down on the intervals  $-5 < x < -3$  and  $0 < x < 2$  because  $g' = f$  is positive and decreasing on these intervals.

2 : { 1 : answer  
1 : reason

(c)  $h'(x) = \frac{5xg'(x) - g(x)5}{(5x)^2} = \frac{5xg'(x) - 5g(x)}{25x^2}$

3 : { 2 :  $h'(x)$   
1 : answer

$$h'(3) = \frac{(5)(3)g'(3) - 5g(3)}{25 \cdot 3^2}$$

$$= \frac{15(-2) - 5(9)}{225} = \frac{-75}{225} = -\frac{1}{3}$$

(d)  $p'(x) = f'(x^2 - x)(2x - 1)$

3 : { 2 :  $p'(x)$   
1 : answer

$$p'(-1) = f'(2)(-3) = (-2)(-3) = 6$$

**AP<sup>®</sup> CALCULUS AB/CALCULUS BC  
2014 SCORING GUIDELINES**

**Question 4**

Train  $A$  runs back and forth on an east-west section of railroad track. Train  $A$ 's velocity, measured in meters per minute, is given by a differentiable function  $v_A(t)$ , where time  $t$  is measured in minutes. Selected values for  $v_A(t)$  are given in the table above.

$t$ (minutes)	0	2	5	8	12
$v_A(t)$ (meters/minute)	0	100	40	-120	-150

- (a) Find the average acceleration of train  $A$  over the interval  $2 \leq t \leq 8$ .
- (b) Do the data in the table support the conclusion that train  $A$ 's velocity is  $-100$  meters per minute at some time  $t$  with  $5 < t < 8$ ? Give a reason for your answer.
- (c) At time  $t = 2$ , train  $A$ 's position is 300 meters east of the Origin Station, and the train is moving to the east. Write an expression involving an integral that gives the position of train  $A$ , in meters from the Origin Station, at time  $t = 12$ . Use a trapezoidal sum with three subintervals indicated by the table to approximate the position of the train at time  $t = 12$ .
- (d) A second train, train  $B$ , travels north from the Origin Station. At time  $t$  the velocity of train  $B$  is given by  $v_B(t) = -5t^2 + 60t + 25$ , and at time  $t = 2$  the train is 400 meters north of the station. Find the rate, in meters per minute, at which the distance between train  $A$  and train  $B$  is changing at time  $t = 2$ .

(a) average accel =  $\frac{v_A(8) - v_A(2)}{8 - 2} = \frac{-120 - 100}{6} = -\frac{110}{3}$  m/min<sup>2</sup>

1 : average acceleration

(b)  $v_A$  is differentiable  $\Rightarrow v_A$  is continuous  
 $v_A(8) = -120 < -100 < 40 = v_A(5)$

2 :  $\begin{cases} 1 : v_A(8) < -100 < v_A(5) \\ 1 : \text{conclusion, using IVT} \end{cases}$

Therefore, by the Intermediate Value Theorem, there is a time  $t$ ,  $5 < t < 8$ , such that  $v_A(t) = -100$ .

(c)  $s_A(12) = s_A(2) + \int_2^{12} v_A(t) dt = 300 + \int_2^{12} v_A(t) dt$   
 $\int_2^{12} v_A(t) dt \approx 3 \cdot \frac{100 + 40}{2} + 3 \cdot \frac{40 - 120}{2} + 4 \cdot \frac{-120 - 150}{2}$   
 $= -450$

3 :  $\begin{cases} 1 : \text{position expression} \\ 1 : \text{trapezoidal sum} \\ 1 : \text{position at time } t = 12 \end{cases}$

$s_A(12) \approx 300 - 450 = -150$

The position of Train  $A$  at time  $t = 12$  minutes is approximately 150 meters west of Origin Station.

- (d) Let  $x$  be train  $A$ 's position,  $y$  train  $B$ 's position, and  $z$  the distance between train  $A$  and train  $B$ .

$z^2 = x^2 + y^2 \Rightarrow 2z \frac{dz}{dt} = 2x \frac{dx}{dt} + 2y \frac{dy}{dt}$

$x = 300, y = 400 \Rightarrow z = 500$

$v_B(2) = -20 + 120 + 25 = 125$

$500 \frac{dz}{dt} = (300)(100) + (400)(125)$

$\frac{dz}{dt} = \frac{80000}{500} = 160$  meters per minute

3 :  $\begin{cases} 2 : \text{implicit differentiation of} \\ \quad \text{distance relationship} \\ 1 : \text{answer} \end{cases}$

**AP<sup>®</sup> CALCULUS AB  
2014 SCORING GUIDELINES**

**Question 5**

$x$	-2	$-2 < x < -1$	-1	$-1 < x < 1$	1	$1 < x < 3$	3
$f(x)$	12	Positive	8	Positive	2	Positive	7
$f'(x)$	-5	Negative	0	Negative	0	Positive	$\frac{1}{2}$
$g(x)$	-1	Negative	0	Positive	3	Positive	1
$g'(x)$	2	Positive	$\frac{3}{2}$	Positive	0	Negative	-2

The twice-differentiable functions  $f$  and  $g$  are defined for all real numbers  $x$ . Values of  $f$ ,  $f'$ ,  $g$ , and  $g'$  for various values of  $x$  are given in the table above.

- (a) Find the  $x$ -coordinate of each relative minimum of  $f$  on the interval  $[-2, 3]$ . Justify your answers.
- (b) Explain why there must be a value  $c$ , for  $-1 < c < 1$ , such that  $f''(c) = 0$ .
- (c) The function  $h$  is defined by  $h(x) = \ln(f(x))$ . Find  $h'(3)$ . Show the computations that lead to your answer.
- (d) Evaluate  $\int_{-2}^3 f'(g(x))g'(x) dx$ .

(a)  $x = 1$  is the only critical point at which  $f'$  changes sign from negative to positive. Therefore,  $f$  has a relative minimum at  $x = 1$ .

(b)  $f'$  is differentiable  $\Rightarrow f'$  is continuous on the interval  $-1 \leq x \leq 1$

$$\frac{f'(1) - f'(-1)}{1 - (-1)} = \frac{0 - 0}{2} = 0$$

Therefore, by the Mean Value Theorem, there is at least one value  $c$ ,  $-1 < c < 1$ , such that  $f''(c) = 0$ .

(c)  $h'(x) = \frac{1}{f(x)} \cdot f'(x)$

$$h'(3) = \frac{1}{f(3)} \cdot f'(3) = \frac{1}{7} \cdot \frac{1}{2} = \frac{1}{14}$$

(d)  $\int_{-2}^3 f'(g(x))g'(x) dx = [f(g(x))]_{x=-2}^{x=3}$   
 $= f(g(3)) - f(g(-2))$   
 $= f(1) - f(-1)$   
 $= 2 - 8 = -6$

1 : answer with justification

$$2 : \begin{cases} 1 : f'(1) - f'(-1) = 0 \\ 1 : \text{explanation, using Mean Value Theorem} \end{cases}$$

$$3 : \begin{cases} 2 : h'(x) \\ 1 : \text{answer} \end{cases}$$


$$3 : \begin{cases} 2 : \text{Fundamental Theorem of Calculus} \\ 1 : \text{answer} \end{cases}$$

**AP<sup>®</sup> CALCULUS AB  
2014 SCORING GUIDELINES**

**Question 6**

Consider the differential equation  $\frac{dy}{dx} = (3 - y)\cos x$ . Let  $y = f(x)$  be the particular solution to the differential equation with the initial condition  $f(0) = 1$ . The function  $f$  is defined for all real numbers.

- (a) A portion of the slope field of the differential equation is given below. Sketch the solution curve through the point  $(0, 1)$ .
- (b) Write an equation for the line tangent to the solution curve in part (a) at the point  $(0, 1)$ . Use the equation to approximate  $f(0.2)$ .
- (c) Find  $y = f(x)$ , the particular solution to the differential equation with the initial condition  $f(0) = 1$ .



1 : solution curve

- (b)  $\left. \frac{dy}{dx} \right|_{(x,y)=(0,1)} = 2 \cos 0 = 2$   
 An equation for the tangent line is  $y = 2x + 1$ .  
 $f(0.2) \approx 2(0.2) + 1 = 1.4$

2 :  $\begin{cases} 1 : \text{tangent line equation} \\ 1 : \text{approximation} \end{cases}$

- (c)  $\frac{dy}{dx} = (3 - y)\cos x$   
 $\int \frac{dy}{3 - y} = \int \cos x \, dx$   
 $-\ln|3 - y| = \sin x + C$   
 $-\ln 2 = \sin 0 + C \Rightarrow C = -\ln 2$   
 $-\ln|3 - y| = \sin x - \ln 2$   
 Because  $y(0) = 1$ ,  $y < 3$ , so  $|3 - y| = 3 - y$   
 $3 - y = 2e^{-\sin x}$   
 $y = 3 - 2e^{-\sin x}$   
 Note: this solution is valid for all real numbers.

6 :  $\begin{cases} 1 : \text{separation of variables} \\ 2 : \text{antiderivatives} \\ 1 : \text{constant of integration} \\ 1 : \text{uses initial condition} \\ 1 : \text{solves for } y \end{cases}$

Note: max 3/6 [1-2-0-0-0] if no constant of integration

Note: 0/6 if no separation of variables