

CAMBRIDGE INTERNATIONAL EXAMINATIONS
Cambridge International General Certificate of Secondary Education

MARK SCHEME for the May/June 2015 series

0606 ADDITIONAL MATHEMATICS

0606/22

Paper 2, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

| | | | |
|--------|---------------------------------|----------|----------|
| Page 2 | Mark Scheme | Syllabus | Paper 22 |
| | Cambridge IGCSE – May/June 2015 | 0606 | |

Abbreviations

| | |
|------|----------------------------|
| awrt | answers which round to |
| cao | correct answer only |
| dep | dependent |
| FT | follow through after error |
| isw | ignore subsequent working |
| oe | or equivalent |
| rot | rounded or truncated |
| SC | Special Case |
| soi | seen or implied |
| www | without wrong working |

| | | | | |
|---|-------|--|----------|--|
| 1 | (i) | | B3,2,1,0 | 2 correctly placed in Venn diagram; 1, 3, 4, 6 correctly placed; 12, 8, 0, 7, 9, 10 correctly placed; 11, 5 correctly placed |
| | (ii) | 3 | B1ft | correct or correct ft <i>their</i> (i), provided non-zero |
| | (iii) | {4, 6} | B1ft | correct or correct ft <i>their</i> (i), provided not the empty set |
| 2 | (i) | $[P =] \begin{pmatrix} 60 & 70 & 58 \\ 50 & 52 & 34 \end{pmatrix}$ and $[Q =] (120 \quad 300)$ | B2 | or $[P =] \begin{pmatrix} 50 & 52 & 34 \\ 60 & 70 & 58 \end{pmatrix}$ and $[Q =] (300 \quad 120)$ or B1 if one error may be written as an unevaluated product; B0 if choice of P and Q offered |
| | (ii) | (22200 24000 17160) | B2 | must have brackets and must not have commas; must be a 1 by 3 matrix; must be from correct product; working may be seen in (i) or B1 for any two elements correct |
| | (iii) | The total (amount of revenue) from all (three) flights. oe | B1 | do not accept, e.g. The total amount from each flight; must be a comment not just a figure; must not contain a contradiction |

| | | | |
|--------|---------------------------------|----------|-------|
| Page 3 | Mark Scheme | Syllabus | Paper |
| | Cambridge IGCSE – May/June 2015 | 0606 | 22 |

| | | | |
|--|--|---|--|
| <p>3 (i)</p> $\frac{(36+15\sqrt{5})}{(6+3\sqrt{5})} \times \frac{(6-3\sqrt{5})}{(6-3\sqrt{5})} \text{ oe}$ $\frac{216+90\sqrt{5}-108\sqrt{5}-225}{-9}$ <p>$1+2\sqrt{5}$ cao</p> <p>Alternative method: $36+15\sqrt{5} = (6a+15b) + (3a+6b)\sqrt{5}$</p> <p>$6a+15b=36$ $3a+6b=15$</p> <p>$a=1$ and $b=2$</p> <p>(ii)</p> $\left[AC^2 = (6+3\sqrt{5})^2 + \text{their } (1+2\sqrt{5})^2 \right]$ $= 36+36\sqrt{5}+45 + \text{their } (1+4\sqrt{5}+20)$ <p>$102+40\sqrt{5}$ cao</p> | | <p>M1</p> <p>DM1</p> <p>A1</p> <p>M1</p> <p>DM1</p> <p>A1</p> <p>M1</p> <p>A1</p> | <p>or $\frac{(12+5\sqrt{5})}{(2+\sqrt{5})} \times \frac{(2-\sqrt{5})}{2-\sqrt{5}}$ oe</p> <p>or $\frac{24+10\sqrt{5}-12\sqrt{5}-25}{-1}$</p> <p>or $-(24+10\sqrt{5})-12\sqrt{5}-25$</p> <p>allow $a=1$ and $b=2$</p> <p>or $1+2\sqrt{5}$</p> <p>correct or correct ft expansions, using Pythagoras with $(6+3\sqrt{5})$ and <i>their</i> BC</p> <p>ignore attempts to square root after correct answer seen</p> |
| <p>4 (i)</p> $\cos(x) = \frac{2}{3} \text{ oe soi}$ <p>$48.189\dots^\circ$ or $131.810\dots^\circ$ or $0.8410\dots$ rad or $2.3(00\dots)$ rad oe isw</p> <p>with reference axis indicated by comment, e.g. “to the bank” or “upstream”, etc. or clearly marked on a diagram</p> | | <p>M1</p> <p>A1</p> | <p>Alternatively</p> $\sin(y) = \frac{2}{3} \text{ oe soi}$ <p>$41.810\dots^\circ$ or $0.7297\dots$ or $0.73(0)$ rad oe isw</p> <p>with reference axis indicated by comment, e.g. “to the perpendicular with the bank”, etc. or clearly marked on a diagram</p> <p>If M0 then SC1 for an unsupported answer of $138.189\dots^\circ$ or $2.4118\dots$ rad or $318.189\dots^\circ$ or $5.5534\dots$ rad with reference axis indicated by comment, e.g. “on a bearing of” or “from North” or clearly marked on a diagram</p> |

| | | | |
|--------|---------------------------------|----------|-------|
| Page 4 | Mark Scheme | Syllabus | Paper |
| | Cambridge IGCSE – May/June 2015 | 0606 | 22 |

| | | | |
|---------------------|--|---|---|
| <p>(ii)</p> | <p>Speed = $\sqrt{9-4}$ ($=\sqrt{5}$) or $3 \sin 48.2$ or $2 \tan 48.2$ or $3 \cos 41.8$ or $\frac{2}{\tan 41.8}$ or $\sqrt{2^2 + 3^2 - 2 \times 2 \times 3 \cos 48.2}$ oe or 2.236(0...) rot to 4 or more figs or 2.24 [m/s] soi</p> <p>time = $\frac{80}{\text{their } \sqrt{5}}$ oe 35.66 to 35.8 (seconds) oe</p> | <p>B1</p> <p>M1</p> <p>A1</p> | <p>Or Distance = $\frac{80}{\sin 48.2} = 107.(33\dots)$ oe soi</p> <p>time = $\frac{\text{their } 107.33\dots}{3}$</p> <p>ignore subsequent rounding or attempted conversion to, e.g. minutes but A0 if answer spoiled by continuation of method</p> <p>if no working, so B0 M0, then allow B3 for an answer 35.66 to 35.8 oe</p> |
| <p>5</p> | <p>Substitution of either $4 - x$ or $4 - y$ into equation of curve and brackets expanded</p> <p>$12x^2 - 52x + 48 [= 0]$ or $12y^2 - 44y + 32 [= 0]$ oe</p> <p>Solve their 3-term quadratic</p> <p>$x = \frac{4}{3}$ and 3 isw</p> <p>$y = \frac{8}{3}$ and 1 isw</p> | <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>A1</p> | <p>condone one sign error or slip in either equation of curve or expansion of brackets; condone omission of $= 0$, BUT $4 - x$ or $4 - y$ must be correct</p> <p>dep on a valid substitution attempt</p> <p>or $x = \frac{4}{3}$ $y = \frac{8}{3}$ not from wrong working</p> <p>or $x = 3$ $y = 1$ not from wrong working</p> <p>if no working, allow full marks for fully correct answer only.</p> |
| <p>6 (a)</p> | <p>$(x-2) \log 6 = \log \left(\frac{1}{4}\right)$ oe or</p> <p>$\log_6 \left(\frac{1}{4}\right) = x-2$ oe</p> <p>1.23 or 1.226(29...) rot to 4 or more figures isw</p> | <p>M1</p> <p>A1</p> | <p>or $x \log 6 = \log \left(\frac{36}{4}\right)$ oe</p> <p>or $x \log 6 - \log 36 = \log 1 - \log 4$ oe</p> <p>correct answer or 1.22 implies M1</p> |

| | | | |
|--------|---------------------------------|----------|-------|
| Page 5 | Mark Scheme | Syllabus | Paper |
| | Cambridge IGCSE – May/June 2015 | 0606 | 22 |


| | | | |
|-----|--|---|---|
| (b) | <p>Method 1</p> $\log\left(\frac{8 \times 2y^2 \times 16y}{64y}\right) = \log 4^2 \text{ oe}$ <p>$y = 2$</p> | <p>B3</p> <p>B1</p> | <p>or B2 if at most one error or omitted step or B1 if at most two errors or omitted steps not from wrong working</p> |
| | <p>Method 2</p> $\log 2 + 2 \log y + 3 \log 2 + 4 \log 2 + \log y - 6 \log 2 - \log y = 4 \log 2$ <p>$y = 2$</p> | <p>B3,2,1,0</p> <p>B1</p> | <p><u>LHS terms</u> $\log 2y^2 = \log 2 + 2 \log y$; $\log 8 = 3 \log 2$; $\log 16y = 4 \log 2 + \log y$; $-\log 64y = -6 \log 2 - \log y$; <u>RHS term</u> $2 \log 4 = 4 \log 2$</p> <p>not from wrong working</p> |
| 7 | $\frac{n(n-1)(n-2)(n-3)(2^4)}{4 \times 3 \times 2 \times 1} = 10 \frac{n(n-1)(2^2)}{2 \times 1}$ <p>or better</p> <p>$n^2 - 5n - 24 [= 0] \text{ oe}$</p> <p>$(n + 3)(n - 8) [= 0]$</p> <p>$n = 8 \text{ only}$</p> | <p>M3</p> <p>A1</p> <p>M1</p> <p>A1</p> | <p>condone omitting the factor of n and/or $n - 1$; must have dealt with factorials</p> <p>M2 if one slip/omission or M1 if two slips/omissions</p> <p>or</p> <p>B1 for $\frac{n(n-1)}{2}(2)^2[x^2]$ seen and B1 for $\frac{n(n-1)(n-2)(n-3)}{24}(2)^4[x^4]$</p> <p>seen equivalent must be 3-terms, e.g. $n^2 - 5n = 24$</p> <p>or any valid method of solution for their 3-term quadratic</p> <p>A0 if -3 also given as a final solution, i.e. not discarded If zero scored, allow SC1 for $n = 8$ unsupported or without correct method</p> |

| | | | |
|--------|---------------------------------|----------|---------|
| Page 6 | Mark Scheme | Syllabus | Page 22 |
| | Cambridge IGCSE – May/June 2015 | 0606 | |

| | | | |
|---|--|--|--|
| 8 | <p>Method 1 (Separate areas subtracted)</p> <p>$[x_B = x_C =] 7$ soi</p> $\left[\int (x^2 - 6x + 10) dx = \right] \frac{x^3}{3} - \frac{6x^2}{2} + 10x$ <p>Correct or correct ft substitution of limits 0 and <i>their</i> 7 into <i>their</i> $\left[\frac{x^3}{3} - \frac{6x^2}{2} + 10x \right]$</p> $\frac{1}{2}(10+17) \times 7 \text{ oe or}$ $\int_0^7 (x+10) dx = \left[\frac{x^2}{2} + 10x \right]_0^7 = \frac{(7)^2}{2} + 10(7) \text{ oe}$ <p><i>their</i> $\left(\frac{189}{2} - \frac{112}{3} \right)$</p> $\frac{343}{6} \text{ or } 57\frac{1}{6} \text{ or } 57.2 \text{ to 3 sf or } 57.16(6\dots)$ <p>rot to 4 figs isw</p> <p>Method 2 (Subtracting and using integration once)</p> <p>$[x_B = x_C =] 7$ soi</p> $\int (-x^2 + 7x) dx$ $\left[-\frac{x^3}{3} + \frac{7x^2}{2} \right] \text{ oe or } \left[\frac{x^3}{3} - \frac{7x^2}{2} \right] \text{ oe}$ <p>Correct or correct ft substitution of limits 0 and <i>their</i> 7</p> <p>into <i>their</i> $\left[-\frac{x^3}{3} + \frac{7x^2}{2} \right]$</p> $\frac{343}{6} \text{ or } 57\frac{1}{6} \text{ or } 57.2 \text{ to 3 sf or } 57.16(6\dots)$ <p>rot to 4 figs isw</p> | <p>B1</p> <p>M2</p> <p>DM1</p> <p>B2</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p> <p>M3</p> <p>M2</p> <p>A1</p> | <p>or M1 for at least one term correct</p> <p>dep on at least M1 being earned; evidence of substitution must be seen in <i>their</i> integral which must be at least two terms; condone omission of lower limit;</p> <p>or M1 for</p> $\frac{1}{2}(\text{their } 10 + \text{their } 17) \times \text{their } 7 \text{ oe}$ <p>or B1 for</p> $\int (x+10) dx = \frac{x^2}{2} + 10x$ <p>dep on a genuine attempt to integrate the equation of the curve; must be <i>their</i> area trapezium/under the line – <i>their</i> attempt at area under curve</p> <p>from full and correct working with no omitted steps</p> <p>condone omission of dx</p> <p>or M2 for</p> $\int (px^2 + qx) dx = \frac{px^3}{3} + \frac{qx^2}{2} \text{ oe either with } p = \pm 1 \text{ or } q = \pm 7$ <p>or M1 for $\int (px^2 + qx) dx = \frac{px^3}{3} + \frac{qx^2}{2}$ with non-zero constants p and q, with $p \neq \pm 1$ and $q \neq \pm 7$</p> <p>dep on a valid integration attempt; evidence of substitution must be seen; condone omission of lower limit;</p> <p>from full and correct working with no omitted steps</p> |
|---|--|--|--|

| | | | | |
|--------|--|--|---|---|
| 9 | (i) | $10 = 2m + 4$ soi $m = 3$ | M1 | or $[m =]\frac{10-4}{2-0}$ oe soi |
| | (ii) | 1 | A1 | |
| | (iii) | $\frac{10 - y_R}{2 - -1} = 1$ oe soi $(-1, 7)$ or $x = -1$ and $y = 7$ | M1 | or $y = x + 8$ oe |
| | | | A1 | if $y = 7$ only stated, provided that $x = -1$ is soi in working allow both marks |
| | | | | if M0 then B1 for $y = 7$ only with no working |
| | (iv) | Use of $m_1 m_2 = -1$ with <i>their</i> m from (i) $y - 10 = \left(\textit{their} - \frac{1}{3}\right)(x - 2)$ $3y + x = 32$ isw | M1 | may be implied by perpendicular gradient seen in equation |
| | | A1 | or $\left(\textit{their} - \frac{1}{3}\right)x + c$ and | |
| | | | $10 = \left(\textit{their} - \frac{1}{3}\right)2 + c$ | |
| | | A1 | allow for correct equation with integer coefficients in any simplified form | |
| (v) | $\left(\frac{1}{2}, \textit{their} \frac{11}{2}\right)$ oe isw | B1, B1ft | ft <i>their</i> y_0 | |
| | | | or M1 for $\left(\frac{2-1}{2}, \frac{10+1}{2}\right)$ seen | |
| (vi) | 4.5 oe cao | B2 | not from wrong working | |
| | | | or M1 for any correct method with correct coordinates | |
| 10 (a) | | B2, 1, 0 | correct sinusoidal/reflected sinusoidal shape, all above x -axis with intent to have all maximum points of equal height; 2 maximum points of intended equal height only over 0 to 360; all max points clearly at $y = 1$; cusp at 180 | |

| | | | |
|--------|---------------------------------|----------|----------|
| Page 8 | Mark Scheme | Syllabus | Paper 22 |
| | Cambridge IGCSE – May/June 2015 | 0606 | |

| | | | | |
|--------|---|--|--|--|
| (b)(i) | $[hg(x) =] \frac{e^{\ln(4x-3)} + 3}{4}$ | M1 | Alternative method $y = \ln(4x - 3)$ and change of subject to x oe, | |
| | fully correct and completion to $[hg(x) =] x$ | A1 | fully correct and comment that $h(x) = g^{-1}(x)$ oe | |
| | (ii) | B2,1,0 | correct shape; 1 marked on the y-axis or (0, 1) stated close by; curve with positive gradient in first quadrant only | |
| |  | | | |
| (iii) | $x \geq 0$ or $[0, \infty)$ | B1 | not domain ≥ 0 | |
| (iv) | $y \geq 1$ or $[1, \infty)$ | B1 | or $h(x) \geq 1$, $h \geq 1$ etc. | |
| 11 | (i) | $\frac{8-h}{8}$ or $8 : 8 - h$ soi | M1 | or $\frac{8}{8-h}$ or $8 - h : 8$ soi |
| | | $\frac{8-h}{8} \times 4$ oe | A1 | or $4 \div \frac{8}{8-h}$ oe |
| | | $h \left(\frac{8-h}{8} \times 4 \right)^2$ oe | M1 | h must be in the numerator of the expression for this mark; |
| | | expand and simplify to $\frac{h^3}{4} - 4h^2 + 16h$ AG | A1 | |
| | (ii) | $\frac{3}{4}h^2 - 8h + 16$ oe | B1 | |
| | | their $\left(\frac{3}{4}h^2 - 8h + 16 \right) = 0$ and attempt to solve | M1 | must be a 3-term quadratic; must be an attempt at a derivative |
| | $\frac{8}{3}$ oe only | A2 | or A1 for $h = \frac{8}{3}$ and 8 allow 2.67 or 2.66(6...) rot to 4 or more figs for $\frac{8}{3}$ | |

| | | | |
|--------|---------------------------------|----------|----------|
| Page 9 | Mark Scheme | Syllabus | Paper 22 |
| | Cambridge IGCSE – May/June 2015 | 0606 | |

| | | | | |
|----|-------|---|--|--|
| 12 | (i) | $-120 + 104 + 22 - 6 = 0$ or correct unsimplified form, e.g. $15(-2)^3 + 26(-2)^2 - 11(-2) - 6 = 0$ or $15(-8) + 26(4) - 11(-2) - 6 = 0$ | B1 | or correct synthetic division $ \begin{array}{r rrrr} -2 & 15 & 26 & -11 & -6 \\ & & -30 & 8 & 6 \\ \hline & 15 & -4 & -3 & 0 \end{array} $ |
| | (ii) | Substituting $x = 3$ into $15x^3 + 26x^2 - 11x - 6$ 600 | M1 | or correct synthetic division $ \begin{array}{r rrrr} 3 & 15 & 26 & -11 & -6 \\ & & 45 & 213 & 606 \\ \hline & 15 & 71 & 202 & 600 \end{array} $ |
| | (iii) | $(x - 1)(15x^3 + 26x^2 - 11x - 6)$ soi Multiply out $(x \pm 1)(15x^3 + 26x^2 - 11x - 6)$ and compare coefficients of x^3 or x to quartic $p = 11$ $q = 5$ | A1 B1 M1 A1 A1 | correct answer implies M1; must be explicitly identified as answer if using synthetic/long division methods by e.g. circling by inspection or division; may be implied by e.g. $(ax + b)(15x^3 + 26x^2 - 11x - 6)$ and $a = 1, b = -1$ seen in later work comparing coefficients or multiply out, e.g. $(ax + b)(15x^3 + 26x^2 - 11x - 6)$ and compare coefficients of x^3 or x to quartic correct p or q implies M1; correct p and q implies B1 M1 |