

**CAMBRIDGE**  
INTERNATIONAL EXAMINATIONS

**NOVEMBER 2002**


**INTERNATIONAL GCSE**

<b>MARK SCHEME</b>
<b>MAXIMUM MARK : 80</b>
<b>SYLLABUS/COMPONENT : 0606/2</b> <b>ADDITIONAL MATHEMATICS</b> <b>(Paper 2)</b>


Page 1	Mark Scheme	Syllabus	Part 2
	IGCSE Examinations – November 2002	0606	



1 [4]	$\text{Inverse} = \begin{pmatrix} 6 & -3 \\ -7 & 4 \end{pmatrix} \times \frac{1}{3}$ $\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 6 & -3 \\ -7 & 4 \end{pmatrix} \begin{pmatrix} -7 \\ -16 \end{pmatrix} = \begin{pmatrix} 2 \\ -5 \end{pmatrix}$	<p>B1 B1</p> <p>M1 A1</p>
2 [4]	$2^6 + 6 \times 2^5 \times x + \frac{6 \times 5}{1 \times 2} \times 2^4 \times x^2$ $= 64 + 192x + 240x^2$ <p>Replace <math>x</math> by <math>x - x^2 \Rightarrow</math> coefficient of <math>x^2 = -192 + 240 = 48</math></p>	<p>B2, 1, 0 (if each incorrect or missing term)</p> <p>M1 A1 c.s.o.</p>
3 [5]	<p>(i) Either <math>\frac{1 + \frac{1}{\sqrt{3}}}{1 - \frac{1}{\sqrt{3}}} \times \frac{1 + \frac{1}{\sqrt{3}}}{1 + \frac{1}{\sqrt{3}}}</math> or <math>\frac{\sqrt{3} + 1}{\sqrt{3} - 1} \times \frac{\sqrt{3} + 1}{\sqrt{3} + 1}</math></p> <p>Simplify <math>\Rightarrow 2 + \sqrt{3}</math></p> <p>(ii) <math>\frac{1}{p} = \frac{1}{2 + \sqrt{3}} \times \frac{2 - \sqrt{3}}{2 - \sqrt{3}} = \frac{2 - \sqrt{3}}{4 - 3}</math></p> $p - \frac{1}{p} = 2 + \sqrt{3} - (2 - \sqrt{3}) = 2\sqrt{3}$ <p>Or <math>p - \frac{1}{p} = 2 + \sqrt{3} - \frac{1}{2 + \sqrt{3}} = \frac{6 + 4\sqrt{3}}{2 + \sqrt{3}}</math></p> <p>Multiply by <math>\frac{2 - \sqrt{3}}{2 - \sqrt{3}} \Rightarrow 2\sqrt{3}</math></p>	<p>M1</p> <p>A1</p> <p>M1 A1 ✓</p> <p>A1</p> <p>B1 ✓</p> <p>M1 A1</p>
4 [6]	<p>Solving inequalities:</p> <p>A <math>x &lt; 3.5</math></p> <p>B <math>x^2 - x - 2 = 0 \Rightarrow (x - 2)(x + 1) = 0 \Rightarrow x = -1, 2</math></p> <p><math>x^2 - x - 2 &gt; 0 \Rightarrow x &lt; -1, x &gt; 2</math></p> <p>Required values <math>-5 &lt; x &lt; -1</math></p> <p><math>2 &lt; x &lt; 3.5</math></p>	<p>B1</p> <p>M1 A1</p> <p>A1</p> <p>M1 A1</p>

Page 2	Mark Scheme	Syllabus	Paper
	IGCSE Examinations – November 2002	0606	2

<p>5 [6]</p>	<p>(a) Either <math>{}_5C_3 = \frac{5 \times 4 \times 3}{1 \times 2 \times 3}</math> or <math>{}_4C_2 = \frac{4 \times 3}{1 \times 2}</math></p> <p>Product = <math>10 \times 6 = 60</math></p> <p>(b) Either, ending in 1 (or 3) <math>\Rightarrow 2 \times 5 \times 4</math> or, ending in 5 (or 7) <math>\Rightarrow 3 \times 5 \times 4</math></p> <p>Adding all 4 cases <math>\Rightarrow 40 + 40 + 60 + 60 = 200</math></p>	<p>B1</p> <p>M1 A1</p> <p>B1</p> <p>M1 A1</p>
<p>6 [6]</p>	<p>(i) <math>f(x) = -(x-1)(x-2)(x-k)</math></p> <p><math>f(3) = -2 \times 1 \times (3-k) = 8 \Rightarrow k = 7</math></p> <p>(ii) <math>f(-3) = -(-4)(-5)(-10) = 200</math></p>	<p>M1 A1</p> <p>M1 A1</p> <p>M1 A1</p>
<p>7 [6]</p>	<p>(i) <math>\frac{d}{dx}(x \sin x) = \sin x + x \cos x</math></p> <p>(ii) <math>\int x \cos x dx = x \sin x - \int \sin x dx</math></p> <p><math>\int \sin x dx = -\cos x</math></p> <p><math>x \sin x + \cos x</math></p> <p><math>\frac{\pi}{2} - 1 \approx 0.571</math></p>	<p>M1 A1</p> <p>M1</p> <p>DM1</p> <p>A1 A1 (e.s.o.)</p>
<p>8 [6]</p>	<p>(i)  [ <math>\rightarrow -\infty</math> as <math>x \rightarrow 0</math>; thro' (1,0); <math>\rightarrow \infty</math> as <math>x \rightarrow \infty</math> ]</p> <p>(ii) Take logs <math>\ln x^2 + \ln e^{x-2} = \ln 1</math></p> <p><math>\Rightarrow 2 \ln x + x - 2 = 0</math></p> <p>Make <math>\ln x</math> the subject <math>\Rightarrow \ln x = -\frac{1}{2}(x-2) \Rightarrow</math> line is <math>y = 1 - x/2</math></p>	<p>B2,1,0</p> <p>M1</p> <p>A1</p> <p>M1 A1</p>
<p>9 [7]</p>	<p>(a) Correct combination of indices</p> <p>Either <math>(a^{2/3} - a^{1/3}b^{2/3} + b^{4/3}) \times a^{1/3} = a - a^{2/3}b^{2/3} + a^{1/3}b^{4/3}</math></p> <p>Or <math>(a^{2/3} - a^{1/3}b^{2/3} + b^{4/3}) \times b^{2/3} = a^{2/3}b^{2/3} - a^{1/3}b^{4/3} + b^2</math></p> <p>Sum = <math>a + b^2</math></p> <p>(b) <math>2^{2x+2} = 4 \times 2^{2x}</math> or <math>2^2 \times 4^x</math></p> <p><math>5^{x-1} = 5^x \div 5</math></p> <p><math>8^x = 2^{3x}</math></p> <p><math>\therefore 10^x = 4/5</math></p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>B2,1,0 (-1 each incorrect or missing term)</p> <p>M1 A1</p>

Page 3	Mark Scheme	Syllabus	0606
	IGCSE Examinations – November 2002		



<p>10 [9]</p>	 <p>(i) <math>AP = \frac{b}{3} - a</math> <span style="margin-left: 200px;"><math>OM = \frac{a}{2} + \frac{b}{2}</math></span></p> <p>(ii) <math>OQ = \lambda (\frac{a}{2} + \frac{b}{2})</math></p> <p>(iii) <math>OQ = OA + \mu AP = a + \mu (\frac{b}{3} - a)</math></p> <p>(iv) Comparing coefficients <math>\lambda/2 = 1 - \mu</math> and <math>\lambda/2 = \mu/3</math></p> <p>Solving <math>\lambda = \frac{1}{2}</math> <span style="margin-left: 50px;"><math>\mu = \frac{2}{3}</math></span></p>	<p>B1 M1 A1</p> <p>B1✓</p> <p>M1 A1✓</p> <p>M1</p> <p>M1 A1</p>
<p>11 [11]</p>	<p>(i) <math>v = \int (\frac{3t}{2} - 6) dt = \frac{3t^2}{4} - 6t (+c)</math></p> <p><math>[v]_{t=0} = 20 \Rightarrow c = 20</math> <span style="margin-left: 200px;"><math>[v]_{t=4} = 12 - 24 + 20 = 8</math></span></p> <p>(ii) <math>\int (\frac{3t^2}{4} - 6t + 20) dt = \frac{t^3}{4} - 3t^2 + 20t</math></p> <p><math>AB = [ ]_0^4 = 16 - 48 + 80 = 48</math></p> <p>(iii) <math>v_B = 8, v_C = 20 \Rightarrow t_{BC} = (20 - 8) / 2 = 6</math></p> <p>(iv) </p> <p style="margin-left: 300px;">curve</p> <p style="margin-left: 300px;">straight line</p>	<p>M1 A1</p> <p>A1 A1</p> <p>M1 A1✓</p> <p>A1</p> <p>M1 A1✓</p> <p>B1</p> <p>B1✓</p>
<p>12 [10]</p> <p>Either</p>	<p><math>A = \pi r^2 + \pi r l \Rightarrow l = (120 - \pi r^2) / \pi r</math></p> <p><math>V = \frac{1}{2} \pi r^2 (\text{expression for } l) = 60r - \frac{1}{2} \pi r^3</math> (AG)</p> <p><math>dV/dr = 60 - 3\pi r^2/2 = 0</math> when <math>r^2 = \frac{40}{\pi} \approx 3.57</math></p> <p>Stationary value of <math>V \approx 143</math> (142.73)</p> <p><math>d^2V/dr^2 = -3\pi r &lt; 0</math> for <math>r &gt; 0 \Rightarrow</math> maximum [or any valid method]</p>	<p>B1 M1</p> <p>M1 A1</p> <p>B1 M1 A1</p> <p>A1</p> <p>M1 A1</p>
<p>Or</p>	<p>(i) <math>dy/dx = x^2 \times 1/x + 2x \ln x</math></p> <p>At Q, <math>y = 0 \Rightarrow \ln x = 0 \Rightarrow x = 1</math> <span style="margin-left: 200px;"><math>[dy/dx]_{x=1} = 1</math> c.s.o.</span></p> <p>(ii) At P, <math>dy/dx = 0 \Rightarrow x(1 + 2 \ln x) = 0 \Rightarrow \ln x = -1/2</math></p> <p><math>\Rightarrow x = e^{-1/2} = 1/\sqrt{e} (\approx 0.6065)</math> (AG)</p> <p>(iii) <math>d^2y/dx^2 = d(x + 2x \ln x) / dx = 1 + 2 \ln x + (2x \times 1/x)</math></p> <p style="margin-left: 200px;"><math>= 3 + 2 \ln x</math></p> <p><math>[d^2y/dx^2]_{x=1/\sqrt{e}} = 3 + 2(-1/2) = 2</math> c.s.o.</p>	<p>M1 A1</p> <p>B1 A1</p> <p>M1 A1</p> <p>A1</p> <p>M1 A1</p> <p>A1</p>