

**Cambridge Assessment International Education** Cambridge International General Certificate of Secondary Education

CANDIDATE NAME							
 CENTRE NUMBER				CANDIDATE NUMBER			
ADDITIONAL MATHEMATICS Paper 2				October/No	0606/21 October/November 2019 2 hours		
Candidates answer on the Question Paper. Additional Materials: Electronic calculator				r			

# **READ THESE INSTRUCTIONS FIRST**

Write your centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, glue or correction fluid. DO NOT WRITE IN ANY BARCODES.

Answer all the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question. The use of an electronic calculator is expected, where appropriate. You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question. The total number of marks for this paper is 80.

www.mymathscloud.com



### Mathematical Formulae

## 1. ALGEBRA

Quadratic Equation

For the equation  $ax^2 + bx + c = 0$ ,

 $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \; .$ 

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \dots + \binom{n}{r}a^{n-r}b^r + \dots + b^n,$$
  
where *n* is a positive integer and  $\binom{n}{r} = \frac{n!}{(n-r)!r!}$ .


### **2. TRIGONOMETRY**

Identities


$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\csc^2 A = 1 + \cot^2 A$$

Formulae for  $\triangle ABC$ 

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$
$$a^2 = b^2 + c^2 - 2bc \cos A$$
$$\Delta = \frac{1}{2} bc \sin A$$







The figure shows part of the graph of  $y = p + q \cos rx$ . Find the value of each of the integers p, q and r.

r =

p = q =

[3]



3 (a) Solve  $e^{2x+1} = 3e^{4-3x}$ .

**(b)** Solve  $\lg(y-6) + \lg(y+15) = 2$ .

[5]

4

#### 4 Do not use a calculator in this question.

www.mymathscloud.com Solve the following simultaneous equations, giving your answers for both x and y in the form  $a+b\sqrt{2}$ , where *a* and *b* are integers.

5

$$2x + y = 5$$
$$3x - \sqrt{2}y = 7$$
[5]

www.mymainscloud.com 5 A particle is moving in a straight line such that t seconds after passing a fixed point O its displacement, str is given by  $s = 3\sin 2t + 4\cos 2t - 4$ .

[3]

[3]


(i) Find expressions for the velocity and acceleration of the particle at time t.

(ii) Find the first time when the particle is instantaneously at rest.

(iii) Find the acceleration of the particle at the time found in part (ii). [2]

#### 6 Do not use a calculator in this question.

www.mymathscloud.com The curve xy = 11x + 5 cuts the line y = x + 10 at the points A and B. The mid-point of AB is the point *C*. Show that the point *C* lies on the line x+y = 11. [7]



(ii) Write p(x) as a product of linear and quadratic factors.

[2]

WWW.MYM3HScloud.com (b) (i) Show that  $13 \tan x \sec x - 4 \sin x - 5 \sec^2 x = 0$  can be written as  $4 \sin^3 x + 9 \sin x - 5 = 0$ .

(ii) Using your answers to part (a)(ii) and part (b)(i) solve the equation


 $13 \tan x \sec x - 4 \sin x - 5 \sec^2 x = 0$  for  $0 < x < 2\pi$  radians. [4]



8 The equation of a curve is given by  $y = xe^{-2x}$ .

(i) Find 
$$\frac{dy}{dx}$$
.

(ii) Find the exact coordinates of the stationary point on the curve  $y = xe^{-2x}$ . [2]



(iv) Using your answer to part (i), find  $\int x e^{-2x} dx$ .

[3]



9 Given that 
$$\mathbf{A} = \begin{pmatrix} 5 & 2 \\ -9 & -3 \end{pmatrix}$$
 and  $\mathbf{B} = \begin{pmatrix} 2 & 1 \\ 6 & 5 \end{pmatrix}$ , find  
(i)  $\mathbf{A}^{-1}$ ,

12

(ii) **B**<sup>2</sup>,

[2]

(iii) the matrix C, where  $B^{-1}C + A = B$ ,

[3]



(iv) the matrix **D**, where  $\mathbf{B}^{-2}\mathbf{D}\mathbf{A} = \mathbf{I}$ .

13



10 (i) Expand  $(3+x)^4$  evaluating each coefficient.

14

In the expansion of  $\left(x - \frac{p}{x}\right)(3+x)^4$  the coefficient of x is zero.

(ii) Find the value of the constant *p*.

(iii) Hence find the term independent of x.

(iv) Show that the coefficient of  $x^2$  is 90.

[2]

[1]

[2]

- www.mymathscloud.com A plane, which can travel at a speed of  $300 \,\mathrm{km}\,\mathrm{h}^{-1}$  in still air, heads due north. The plane is blown off cours by a wind so that it travels on a bearing of  $010^\circ$  at a speed of  $280 \,\mathrm{km}\,\mathrm{h}^{-1}$ . 11
  - (i) Find the speed of the wind.

[3]

(ii) Find the direction of the wind as a bearing correct to the nearest degree.



16

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.