



Cambridge IGCSE™

ADDITIONAL MATHEMATICS

0606/11

Paper 1

October/November 2020

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2020 series for most Cambridge IGCSE™, Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

This document consists of **9** printed pages.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always **whole marks** (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Maths-Specific Marking Principles	
1	Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
2	Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
3	Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
4	Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
5	Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread.
6	Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.


Types of mark

- M** Method marks, awarded for a valid method applied to the problem.
- A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.
- B** Mark for a correct result or statement independent of Method marks.


When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation '**dep**' is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Abbreviations

awrt	answers which round to
cao	correct answer only
dep	dependent
FT	follow through after error
isw	ignore subsequent working
nfww	not from wrong working
oe	or equivalent
rot	rounded or truncated
SC	Special Case
soi	seen or implied

Question	Answer	Marks	Guidance
1	$y = \pm 3(x+2)(x+1)(x-4)$	3	B1 for 3 B1 for $(x+2)(x+1)(x-4)$ B1 for \pm
2(a)	4	B1	
2(b)	1080° or 6π	B1	
2(c)		3	B1 for shape, it must be symmetrical about the y -axis. B1 for y -intercept of 5 B1 for $(\pm 180^\circ, 3)$
3(a)	$a = \frac{3}{2}$ or $p^{\frac{3}{2}}$	B1	
	$b = \frac{10}{3}$ or $q^{\frac{10}{3}}$	B1	
	$c = -\frac{7}{3}$ or $r^{\frac{7}{3}}$	B1	
3(b)	$\left(3x^{\frac{1}{3}} - 1\right)\left(2x^{\frac{1}{3}} - 1\right) = 0$	M1	For recognising as a quadratic in $x^{\frac{1}{3}}$ and attempt to solve to obtain $x^{\frac{1}{3}} = k$
	$x^{\frac{1}{3}} = \frac{1}{3}, x^{\frac{1}{3}} = \frac{1}{2}$ leading to $x = \frac{1}{27}$ or 0.0370 $x = \frac{1}{8}$ or 0.125	2	Dep M1 for a valid method of solving $x^{\frac{1}{3}} = k$ where $k > 0$ A1 for both
4(a)	$\frac{dy}{dx} = \frac{\sin x \times 3\sec^2 3x - \tan 3x \cos x}{\sin^2 x}$	3	B1 for $3\sec^2 3x$ M1 for differentiation of a quotient or equivalent product A1 for all other terms apart from $3\sec^2 3x$ correct
	When $x = \frac{\pi}{3}$ $\frac{dy}{dx} = 2\sqrt{3}$	A1	
4(b)	$2\sqrt{3}h$	B1	FT on <i>their</i> answer to (a)

Question	Answer	Marks	Guidance
4(c)	$\frac{dy}{dx} \times \frac{dx}{dt} = \frac{dy}{dt}$ $2\sqrt{3} \times 3 = \frac{dy}{dt}$	M1	For correct use of rates of change using <i>their</i> answer to (a)
	$\frac{dy}{dt} = 6\sqrt{3}$	A1	
5(a)(i)	360	B1	
5(a)(ii)	Starts with 6: $1 \times 4 \times 3 \times 1 = 12$	B1	
	Starts with 7 or 9 : $= 2 \times 4 \times 3 \times 2 = 48$	B1	
	Total = 60	B1	
	Alternative		
	Ending in 4: $\frac{1}{6} \times 360 \times \frac{3}{5} = 36$	(B1)	Allow unsimplified
	Ending in 6: $\frac{1}{6} \times 360 \times \frac{2}{5} = 24$	(B1)	Allow unsimplified
	Total = 60	(B1)	
5(b)(i)	1287	B1	
5(b)(ii)	$1287 - {}^7C_5$ or 1 doctor: 210 2 doctors: 525 3 doctors: 420 4 doctors: 105 5 doctors: 1	M1	For <i>their</i> (b)(i) $-{}^7C_5$ or listing all the other separate cases which must be evaluated, allow 1 error
	1266	A1	
5(b)(iii)	45	B1	
6(a)	Velocity vector = $\begin{pmatrix} -8 \\ 6 \end{pmatrix}$	2	M1 for obtaining 5
	$\begin{pmatrix} 30 \\ 10 \end{pmatrix} + \begin{pmatrix} -8 \\ 6 \end{pmatrix} t$	B1	FT for $\begin{pmatrix} 30 \\ 10 \end{pmatrix} + (\textit{their} \text{ velocity vector})t$
6(b)	13	B1	

Question	Answer	Marks	Guidance
6(c)	$P: \begin{pmatrix} -50 \\ 70 \end{pmatrix}$ $Q: \begin{pmatrix} -30 \\ 210 \end{pmatrix}$	M1	Using $t = 10$ to find position vector of each particle
	$\sqrt{20^2 + 140^2}$	M1	Dep on previous M mark, for use of Pythagoras on difference of the 2 position vectors
	$100\sqrt{2}$	A1	
7(a)	$f \in \mathbb{R}$	B1	Allow y but not x
7(b)	$x = 5 \ln(2y + 3)$ $e^{\frac{x}{5}} = 2y + 3$	M1	For a complete attempt to obtain inverse
	$f^{-1}(x) = \frac{e^{\frac{x}{5}} - 3}{2}$	A1	Must be using correct notation
	Domain $x \in \mathbb{R}$	B1	FT on <i>their</i> (a). Must be using correct notation
7(c)		5	B1 for shape of $y = f(x)$ B1 for shape of $y = f^{-1}(x)$ B1 for 5 ln 3 or 5.5 and -1 on both axes for $y = f(x)$ B1 for 5 ln 3 or 5.5 and -1 on both axes for $y = f^{-1}(x)$ B1 All correct, with apparent symmetry which may be implied by previous 2 B marks or by inclusion of $y = x$, and implied asymptotes, may have one or two points of intersection
8(a)(i)	$\frac{1}{\left(1 + \frac{1}{\sin \theta}\right)(\sin \theta - \sin^2 \theta)}$	B1	For use of $\operatorname{cosec} \theta = \frac{1}{\sin \theta}$, may be implied
	$\frac{1}{\sin \theta + 1 - \sin \theta - \sin^2 \theta}$	M1	For expansion of brackets
	$\frac{1}{\cos^2 \theta}$	M1	For simplification and use of identity
	$\sec^2 \theta$	A1	For final result, must see $\frac{1}{\cos^2 \theta}$

Question	Answer	Marks	Guidance
8(a)(ii)	$\cos^2 \theta = \frac{3}{4}$	B1	For relating to and making use of (a)
	$\cos \theta = \pm \frac{\sqrt{3}}{2}$	M1	For attempt to solve, may be implied by one correct solution
	$\theta = -150^\circ, -30^\circ, 30^\circ, 150^\circ$	2	A1 for any correct pair A1 for a second correct pair and no extra solutions within the range
8(b)	$\tan\left(3\phi + \frac{2\pi}{3}\right) = 1$	B1	
	$3\phi + \frac{2\pi}{3} = \frac{\pi}{4}, \frac{5\pi}{4}, \frac{9\pi}{4}$ $3\phi = \frac{7\pi}{12}, \frac{19\pi}{12}$	M1	For correct order of operations
	$\phi = \frac{7\pi}{36}$	A1	
	$\phi = \frac{19\pi}{36}$	A1	
9(a)	$\left[\ln x - \frac{1}{2}\ln(2x+3)\right]_1^a$	2	B1 for $\ln x$ B1 for $\frac{1}{2}\ln(2x+3)$
	$\ln a - \frac{1}{2}\ln(2a+3) + \frac{1}{2}\ln 5$	M1	For correct application of limits, must have at least one B1
	$\ln a \sqrt{\frac{5}{2a+3}}$	M1	Dep on previous M mark, for application of log laws
	$5a^2 - 18a - 27 = 0$	M1	Dep on previous M mark for equating to $\ln 3$ and simplification to a 3 term quadratic = 0
	$a = \frac{9+6\sqrt{6}}{5}$	A1	Must have one solution only

Question	Answer	Marks	Guidance
9(b)	$-\frac{1}{2}\cos\left(2x + \frac{\pi}{3}\right) + \frac{1}{2}\sin 2x - x$	3	B1 for $-\frac{1}{2}\cos\left(2x + \frac{\pi}{3}\right)$ B1 for $+\frac{1}{2}\sin 2x$ B1 for $-x$
	$\left(-\frac{1}{2}\cos\pi + \frac{1}{2}\sin\frac{2\pi}{3} - \frac{\pi}{3}\right)$ $-\left(-\frac{1}{2}\cos\frac{\pi}{3}\right)$	M1	For correct use of limits in <i>their</i> integral, must have at least one B1 term
	$\frac{3}{4} + \frac{\sqrt{3}}{4} - \frac{\pi}{3}$	A1	
10(a)	$a + d = 8$ $a + 3d = 18$	2	B1 for both equations M1 for attempt to solve <i>their</i> equations
	$a = 3, d = 5$	A1	For both
	$\frac{n}{2}(6 + (n-1)5) > 1560$	M1	For correct use of sum formula with <i>their</i> a and d , allow equality
	$5n^2 + n - 3120 > 0$	M1	For attempt to solve, allow equality, to obtain at least one critical value
	Positive critical value 24.9 25terms	A1	
10(b)(i)	$\frac{a}{1-r} = 72$ and either $a + ar + ar^2 = \frac{333}{8}$ or $\frac{a(1-r^3)}{1-r} = \frac{333}{8}$	B1	For both
	$a = 72(1-r)$ and $a(1+r+r^2) = \frac{333}{8}$ oe $72(1-r)(1+r+r^2) = \frac{333}{8}$ or $72(1-r^3) = \frac{333}{8}$	M1	For attempt to obtain an equation in terms of r only
	$1-r^3 = \frac{333}{576}$	A1	
	$r = 0.75$	2	M1 for attempt to solve <i>their</i> equation in r

Question	Answer	Marks	Guidance
10(b)(ii)	$a = 18$	B1	FT on their r provided $ r < 1$