



Cambridge International Examinations

Cambridge International General Certificate of Secondary Education

MATHEMATICS
Paper 2 (Extended)
MARK SCHEME
Maximum Mark: 70

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

® IGCSE is a registered trademark.



May, Mynainscloud.com

Abbreviations

cao correct answer only

dep dependent

FT follow through after error isw ignore subsequent working

oe or equivalent SC Special Case

nfww not from wrong working

soi seen or implied

Question	Answer	Marks	Part Marks
1	0.407 or 0.4067	1	
2	4x(x-2y) final answer	2	M1 for $4(x^2 - 2xy)$ or $x(4x - 8y)$ or $2(2x^2 - 4xy)$ or $2x(2x - 4y)$
3	120	2	M1 for finding a correct product of prime factors or correctly listing a minimum of 3 multiples of 20 and 24 or for answer $2^3 \times 3 \times 5$ oe or $120k$ where k is an integer > 1
4	$(x-y)^2$ oe final answer	2	M1 for $x - y = \sqrt{a}$ or their $(x - y)$ squared
5	68.6 or 68.62 to 68.64	2	M1 for $\frac{1}{2} \times \frac{4}{3} \pi \times 3.2^3$ If zero scored, SC1 for final answer 137 or 137.2 to 137.3
6	$\frac{4}{25}$ oe	2	M1 for $\frac{2}{5} \times \frac{2}{5}$ oe or denominator 5^2 oe
7	$\frac{32}{x^2}$ or $32x^{-2}$ final answer	2	M1 for $y = \frac{k}{x^2}$ oe or $[k =]32$
8	$\frac{2}{a^4}$ or $2a^{-4}$ final answer	2	B1 for $\frac{2}{a^k}$ oe or $\frac{k}{a^4}$ oe $(k \neq 0)$ final answer
9(a)(i)	$\begin{pmatrix} 30 \\ -20 \end{pmatrix}$	1	
9(a)(ii)	$\begin{pmatrix} -6 \\ 4 \end{pmatrix}$	1	
9(b)	-4	1	

Question	Answer	Marks	Part Marks
10(a)	10	2	M1 for $5x + 6x + 7x = 180$ oe or $\frac{180}{5 + 6 + 7}$ or B1 for angles 50, 60 and 70
10(b)	70	1FT	FT $7 \times their$ (a) provided $0 < their answer < 180$
11	Correct region	3	1 2 2 1 2 SC1 for R not marked and reverse shading
12(a)	3+12x final answer	1	
12(b)	24x + 31 final answer	2	M1 for $3 + 4(6x + 7)$
13	150	3	M2 for $\left(\frac{1}{0.512}\right)^{\frac{2}{3}}$ oe or $\left(\frac{0.512}{1}\right)^{\frac{2}{3}}$ oe or M1 for scale factor $\left(\frac{1}{0.512}\right)^{\frac{1}{3}}$ oe or $\left(\frac{0.512}{[1]}\right)^{\frac{1}{3}}$ oe
14	$10^{k+2} \times [0].\dot{6}\dot{3} - 10^k \times [0].\dot{6}\dot{3}$ oe where $k > 1$	M1	
	$\frac{63}{99}$ or equivalent fraction	A1	e.g. $\frac{6300}{9900}$ but not $\frac{7}{11}$
	7 11	B1	
15	35.8 or 35.77	3	M2 for $[\sin =] \frac{24 \times \sin 71.8}{39}$ or M1 for $\frac{39}{\sin 71.8} = \frac{24}{\sin x}$ oe
16(a)	$x \le 3$ final answer	2	M1 for $13 - 7 \ge 3x - x$ oe
16(b)	1, 2, 3	1FT	correct answer or FT their answer to (a)

Question	Answer	Marks	Part Marks
17	$\frac{2}{7}\mathbf{p} + \frac{5}{7}\mathbf{q}$	3	M1 for $PZ = \frac{5}{7} (\mathbf{q} - \mathbf{p})$ oe or $QZ = \frac{2}{7} (\mathbf{p} - \mathbf{q})$ oe M1 for correct route from O to Z or identifying OZ
18	3000	3	M2 for $12.5 \times \frac{1}{2} (200 + 280)$ oe or M1 for part area
19	common denominator 12	B1	accept $k \times 12$ throughout
	one correct from $\frac{9}{12}$ or $\frac{8}{12}$ oe	M1	accept $\frac{9k}{12k}$ or $\frac{8k}{12k}$
	$\frac{5}{6}$ cao	A2	A1 for $\frac{10}{12}$ or $\frac{10k}{12k}$
20(a)	6	1	
20(b)	$2x^3$ final answer	1	
20(c)	$15y^4$ final answer	2	B1 for $15y^k$ or ky^4 as final answer $(k \neq 0)$
21	$\sqrt{10^2 - 4 \times 5 \times 2}$ oe or better	B1	If completing the square: B1 for $(x+1)^2$ oe B1 for $-1+\sqrt{1-\frac{2}{5}}$ or $-1-\sqrt{1-\frac{2}{5}}$ oe
	$\frac{-10 + \sqrt{q}}{2(5)}$ or $\frac{-10 - \sqrt{q}}{2(5)}$ oe	B1	
	- 0.23, -1.77 final ans cao	B1B1	SC1 for
			- 0.2 or - 0.225 and -1.8 or -1.774 or -1.775 or 0.23 and 1.77 as answer or - 0.23 and -1.77 seen in working Maximum score without working is 2
22	35.3 or 35.26	4	M3 for $[\tan =] \frac{26}{\sqrt{26^2 + 26^2}}$ oe
			or M1 for $[AC^2 =] 26^2 + 26^2$ oe and M1 for $[\tan =] 26 \div their AC$ oe or for angle CAG indicated

Question	Answer	Marks	Part Marks
23(a)	4(x-6) or $4x-24$ as final answer	1	
23(b)	$x^2 + 23x + 26$ final answer	3	B2 for $x^2 + 4x + 4x + 16$ or better or B1 for $15x + 10$
24	1.96 cao	5	M4 for $\left(\left(\sqrt[3]{\frac{2500 \times 1.6 \times 3}{100} + 2000}\right) - 1\right) - 1\right)$ 100] oe or 1.96or [0].0196 or 101.96 or 1.0196 or 1.0196 or $\sqrt[3]{\frac{2500 \times 1.6 \times 3}{100} + 2000}$ or B2 for [SI =] 120 or [CI total=] 2120 or M1 for $\sqrt[2500 \times 3 \times 1.6}{100}$ and M1 for $2000 \times (k)^3$