
CAMBRIDGE INTERNATIONAL MATHEMATICS

0607/61

Paper 6 (Extended)

May/June 2016

MARK SCHEME

Maximum Mark: 40

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2016 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

Page 2	Mark Scheme	Syllabus	Page
	Cambridge IGCSE – May/June 2016	0607	61

Abbreviations

- cao correct answer only
- dep dependent
- FT follow through after error
- isw ignore subsequent working
- oe or equivalent
- SC Special Case
- nfww not from wrong working
- soi seen or implied

A INVESTIGATION		MOVING TRIANGLES													
Question	Answer	Mark	Part Marks												
1 (a)	2	1	B1 for each one correct												
(b)	<table border="1"> <tr> <th>Scale factor</th> <th>PS</th> <th>PB</th> </tr> <tr> <td>3</td> <td>4</td> <td>12</td> </tr> <tr> <td>5</td> <td>6</td> <td>30</td> </tr> <tr> <td>7</td> <td>2</td> <td>14</td> </tr> </table>	Scale factor		PS	PB	3	4	12	5	6	30	7	2	14	3
Scale factor	PS	PB													
3	4	12													
5	6	30													
7	2	14													
(c)	Similar	1													
2 (a)	$\frac{2}{20} = \frac{1}{10}$ oe	1	Allow, for example, 2 : 20 = 1 : 10 or 2 : 1 = 20 : 10 or 2 × 10 = 20 and 1 × 10 = 10 or 2 : 20 and 1 : x so 2x = 20, x = 10 or PS is double RS so PB is double QB or equivalent												
(b)	8	1	C opportunity												
(c)	$\frac{y}{2}$ oe	1	condone $\frac{y}{2} \times 1$												
3	$\frac{y}{4}$ oe	1	condone $\frac{y}{4} \times 1$ If 0 scored in 2(c) and 3, allow SC1 for answers of $y = 2x$ and $y = 4x$												

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2016	0607	61

Question	Answer	Mark	Part Marks
4 (a)	18	1	C opportunity
(b)	12	1	C opportunity
(c)	<i>their</i> 6	1FT	strict FT <i>their</i> $y - z$
5	$[y =] 5x$ and $[z =] 4x$ $[AP =] 5x - 4x = x$	M1 A1	may be on diagram Allow 2 marks for $y = 5x$ and $z = 4x$ seen or clearly indicated $[AP =] y - z = x$
6	$[AP =] nx - (n - 1)x = x$	1	or $nx - (nx - x) = x$ or $nx - nx + x = x$ not from wrong working or equating expressions for BQ $\frac{y}{n} = \frac{z}{n-1}$ and rearranging to show that either $y - z = \frac{y}{n}$ with $x = \frac{y}{n}$ or that $y - z = \frac{z}{n-1}$ with $x = \frac{z}{n-1}$ C opportunity
7 (a)	$\frac{x}{2}$	2	M1 for $\frac{1}{2}xn$ and $\frac{1}{2}x(n-1)$ oe seen or for $x = 2AP$
(b)	$\frac{x}{m}$	1	C opportunity
Communication seen in 3 of 2(b), 4(a), 4(b), 6 or 7(b)		2	C1 if seen in two of them

B MODELLING		MUSICAL NOTES	
Question	Answer	Mark	Part Marks
1	<p>Correct curve over full domain.</p>	2	<p>B1 for at least one correct, complete cycle e.g. over the domain $0 \leq t \leq \frac{1}{110}$</p> <p>or for a graph of incorrect shape but that has 4 cycles over the full domain</p> <p>or for a graph with more than 3 inaccurate t-intercepts with 4 cycles over the full domain</p> <p>or for a fully correct and accurate sketch graph of the sine wave for the note A_0</p>
2	<p>(a) (i) 32.7[0] or 32.703 to 32.7032 isw</p> <p>(ii) C_1</p> <p>(iii) 41.2[0] or 41.203 to 41.2035 isw</p> <p>(b) [0, 12,] 24, 36, 48, 60, 72, 84</p> <p>(c) C_7 and 4190 or 4186 or 4186.0 or 4186.00 or 4186.009 to 4186.01</p>	1 1 1 1 1	C opportunity
3	$2^{\frac{1}{12}}$ or exact equivalent	1	isw conversion to decimal, but decimal answer only does not score C opportunity
4	<p>(a) Correct exponential shape</p> <p>(b) F_5</p>	1 2	<p>Intent of smooth curve; must not cross x-axis; condone graph not drawn on full domain; condone y-intercept at origin;</p> <p>M1 for $n = 68$ soi e.g. $f(68)$ or $27.5 \times 2^{\frac{68}{12}}$</p> <p>C opportunity</p>

Question	Answer	Mark	Part Marks
5 (a)	600	1	
(b)	$\frac{1}{10}$ oe isw	1	
(c)	Uses an algebraic process to find either $h(n+1) = 2^{\text{their } \frac{1}{10}} \times h(n)$ oe or $k = 2^{\text{their } \frac{1}{10}}$ or 1.07 or 1.071 to 1.072	1FT	FT <i>their</i> value of b , provided $b \neq 1$; Allow $k = 2^b$ isw Condone k found by calculating the ratio of at least 2 pairs of consecutive values e.g. $\frac{h(2)}{h(1)}$ and $\frac{h(4)}{h(3)}$
6 (a)	77.3 or 77.29 to 77.295	2	M1 for $2^{\frac{k}{23}}$ where k may be a constant or a variable seen C opportunity
(b)	9	2	not from wrong working M1 for $100 \times 2^n = 108$ or $100 \times 1.08^n = 200$ or $1.08^n = 2$ or for $1.08^9 = 1.99\dots$ soi or for two correct trials using a valid relationship seen C opportunity
Communication in 2 of 2(a)(iii), 3, 4(b), 6(a) or 6(b)		2	C1 if seen in 1 of them