CANDI	International Ge	OF CAMBRIDGE INTE eneral Certificate of Se	RNATIONAL EXAMINAT	Mun MR	Ud.com
CENTF NUMBI			CANDIDATE NUMBER		
	RIDGE INTERNATIONAL	. MATHEMATICS		0607/02	
Paper 2	! (Extended)		0	ctober/November 2013 45 minutes	
Candid	ates answer on the Quest	ion Paper.			
Additio	nal Materials: Geome	etrical Instruments			

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

Do not use staples, paper clips, highlighters, glue or correction fluid.

You may use a pencil for any diagrams or graphs.

DO NOT WRITE IN ANY BARCODES.

Answer all the questions.

CALCULATORS MUST NOT BE USED IN THIS PAPER.

All answers should be given in their simplest form.

You must show all the relevant working to gain full marks and you will be given marks for correct methods even if your answer is incorrect.


The number of marks is given in brackets [] at the end of each question or part question.

The total number of marks for this paper is 40.



Formula List

For the equation	$ax^2 + bx + c = 0$	$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
Curved surface area, A, of cyli	nder of radius r, height h.	$A = 2\pi rh$
Curved surface area, A, of cond	e of radius r, sloping edge l.	$A = \pi r l$
Curved surface area, A, of sphe	ere of radius <i>r</i> .	$A = 4\pi r^2$
Volume, <i>V</i> , of pyramid, base a	rea A, height h.	$V=\frac{1}{3}Ah$
Volume, V, of cylinder of radiu	us r , height h .	$V = \pi r^2 h$
Volume, V , of cone of radius r	, height <i>h</i> .	$V = \frac{1}{3}\pi r^2 h$
Volume, <i>V</i> , of sphere of radius	r.	$V = \frac{4}{3}\pi r^3$
A		$\underline{a} = \underline{b}$





 $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$ $a^2 = b^2 + c^2 - 2bc \cos A$ $\operatorname{Area} = \frac{1}{2}bc \sin A$

www.mymathscioud.com


© UCLES 2013


2






Sin	plify the following.	
(a)	5 aplify the following. $2y^2 \times 3y^3$	S.S.C
	Answer(a)	[2]
(b)	$\sqrt[3]{27p^{27}}$	
	Answer(b)	[2]
(a)	Find the amplitude and period of the function $f(x) = 4\cos(4x)$.	
	Answer(a) Amplitude =	
	Period =	[2]
(b)	$g(x) = 4\cos(4x) - 4$	
	Describe fully the single transformation that maps the graph of $y = f(x)$ onto the graph $y = g(x)$.	of
	Answer(b)	
		[2]





Question 12 is printed on the next page.



Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.