



Cambridge IGCSE™

CAMBRIDGE INTERNATIONAL MATHEMATICS

0607/42 October/November 2020

Paper 4 (Extended) MARK SCHEME Maximum Mark: 120

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2020 series for most Cambridge IGCSE[™], Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

Generic Marking Principles

October/Nove. W. Mymathscioud.com These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given • for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6[•]

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

| 0607 | 42 Cambridge IGCSE – Mark Scheme October/Nove. Dynam PUBLISHED October/Nove. Dynam | | | | |
|------|---|--|--|--|--|
| Ma | hs-Specific Marking Principles | | | | |
| 1 | Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing. | | | | |
| 2 | Unless specified in the question, answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected. | | | | |
| 3 | Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points. | | | | |
| 4 | Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw). | | | | |
| 5 | Where a candidate has misread a number in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 mark for the misread. | | | | |
| 6 | Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear. | | | | |

MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- Μ Method marks, awarded for a valid method applied to the problem.
- А Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.
- В Mark for a correct result or statement independent of Method marks.

When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation 'dep' is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Abbreviations

- awrt answers which round to
- correct answer only cao
- dep dependent
- FT follow through after error
- ignore subsequent working isw
- not from wrong working nfww
- or equivalent oe
- rounded or truncated rot
- SC Special Case
- seen or implied soi

|)7/42 | Cambridge IGCSE – Mark Scheme Det Det Scheme Det Det Scheme Det Det Scheme Det Det Scheme Det Scheme Det Scheme Det Det Scheme Det Det Scheme Det Scheme Det Det Scheme D | | | |
|----------|---|-------|--|--|
| Question | Answer | Marks | Partial Marks | |
| 1(a) | 23 500 | 2 | M1 for $x \times \frac{100 - 15}{100} = 19975$ oe or better | |
| 1(b)(i) | 22697.5[0] final answer | 3 | M1 for 19975 × $\frac{10}{100}$ soi by 1997.5 M1 for 12 × 345 [× 5] | |
| 1(b)(ii) | 97.62 | 4 | M2 for $19975 \left(1 + \frac{25}{100}\right)^5$ or M1 for $19975 \left(1 + \frac{25}{100}\right)^n$, $n > 1$ M1 for their 22 697.5 – their 22 599.88 | |
| 2(a) | Translation $ \begin{pmatrix} 4 \\ 0 \end{pmatrix} $ | 2 | B1 for each | |
| 2(b) | Triangle at (-1, -1), (-2, -1), (-1, -3) | 2 | B1 for reflection in $y = x$ | |
| 2(c) | Triangle at (3, -1), (4, -1), (3, -3) | 2 | B1 for rotation 90° clockwise, wrong centre or rotation 90° anti-clockwise about (1, -1) | |
| 2(d) | Reflection $x = 1$ | 2 | B1 for each | |
| 2(e) | Stretch x-axis invariant oe [factor] 2 | 3 | B1 for each | |
| 3(a) | 4 correct points | 2 | B1 for 2 or 3 correct points | |
| 3(b) | Negative | 1 | | |
| 3(c) | [y =] -2.84x + 19[.0] | 2 | 2.844, 19.03 B1 for -2.84 <i>x</i> + <i>b</i> or <i>ax</i> +19[.0], <i>a</i> < 0 or -2.8 <i>x</i> +19 | |
| 3(d) | 11[.0] or 11.1 or 11.04 to 11.08 | 1 | FT their (c) | |

| 07/42 | Mark Scheme October/Nove. Munimitient Cambridge IGCSE – Mark Scheme October/Nove. Output Output <th< th=""></th<> | | | |
|----------|---|-------|--|--|
| Question | Answer | Marks | Partial Marks | |
| 4(a) | Correct sketch | 2 | Cubic curve, crossing <i>x</i> -axis three times, with maximum in 2nd quadrant and minimum, in 4th quadrant B1 for any cubic (positive <i>x</i> ³) | |
| 4(b) | (1.29, -1.3[0]) | 2 | 1.290 to 1.291, -1.303 B1 for each coordinate | |
| 4(c) | Rotation [Order] 2 [centre] (0, 3) | 3 | B1 for each | |
| 4(d)(i) | -2.9[0] or -2.895 0.603 or 0.6027 2.29 or 2.292 | 3 | B1 for each If three correct but seen in coordinate form, SC1 . If 0 scored, SC1 for correct sketch of line (may be freehand but negative intersection with <i>y</i> -axis and crosses curve three times) | |
| 4(d)(ii) | -2.9[0] < x < 0.603 x > 2.29 | 2 | FT <i>their</i> (i) B1 for each | |
| 5(a) | 866 or 865.5 to 865.6 | 2 | M1 for 14498 ÷ 16.75 | |
| 5(b) | 1400 or 2 pm | 3 | B1 for 29h 60min or 30 h B1 for 06 00 | |
| | | | OR | |
| | | | B1 for 21 15 oe B1 for 38h or 37h 60min | |
| | | | OR | |
| | | | M1 for 13 15 + 8 + 16 45 or 13 15 + 16 45 + 8 M1 for a correct conversion to 24 hour clock | |
| 5(c) | 1505 cao | 2 | M1 for 827.75 ÷ 0.55 | |

| 07/42 | | IGCSE – Mark Scl PUBLISHED | heme October/Nove. Multiply $Partial Marks$ M1 for $62^2 = 44^2 + 95^2 - 2 \times 44 \times$ |
|----------|--|-------------------------------|---|
| Question | Answer | Marks | Partial Marks |
| 6(a) | $[\cos =]\frac{44^2 + 95^2 - 62^2}{2 \times 44 \times 95}$ | M2 | M1 for $62^2 = 44^2 + 95^2 - 2 \times 44 \times 95\cos[]$ |
| | 31.64 | A1 | |
| 6(b) | 145 or 145.4 to 145.5 | 3 | B2 for [angle $DAC =$] 38.8 or 38.83 to 38.84 or M1 for cos [DAC] = $\frac{74}{95}$ oe |
| 6(c) | 23.1 or 23.05 to 23.09 | 2 | M1 for $\frac{distance}{44} = \sin 31.6$ oe |
| 6(d) | 3300 or 3296 to 3302 | 3 | M1 for $0.5 \times 95 \times their(c)$ oe M1 for $0.5 \times 74 \times 95 \times sin(their 38.8)$ oe |
| 7(a) | 6.71 or 6.708 | 3 | M2 for $(9-3)^2 + (5-2)^2$ soi or M1 for $9-3$ and $5-2$ soi |
| 7(b) | [y=] 0.5x + 0.5 oe | 3 | M1 for gradient = $\frac{5-2}{9-3}$ oe M1 for substituting (3, 2) or (9, 5) into <i>their</i> $y = mx + c$ oe |
| 7(c) | [y=]-2x+18 | 3 | M1 for gradient = $-1 \div their 0.5$ M1 for substituting (8, 2) into <i>their</i> y = mx + c oe |
| 7(d) | (7, 4) | 2 | B1 for each coordinate or M1 for <i>their</i> (b) = <i>their</i> (c) |
| 7(e)(i) | (6, 6) | 2 | B1 for each coordinate |
| 7(e)(ii) | Kite | 1 | |

| 0607/42 | Cambridge IGCSE PUBLIS | | heme October/Nove. My have $Partial Marks$ M2 for $\frac{(8-3)\times(5-2)}{2}\times 2$ | |
|----------|---------------------------------------|-------|--|-----|
| Question | Answer | Marks | Partial Marks | |
| 7(f) | 15 or 14.99 to 15.01 | 3 | 2 | -ON |
| | | | OR M1 for $((their 6) - 8)^2 + ((their 6) - 2)^2$ M1 for $0.5 \times their$ (a) $\times their$ CD | |
| | | | OR | |
| | | | M2 for $24 - 6 - 2 \times 1.5$ oe or better or M1 for $6 \times 4 - 0.5 \times 3 \times 4 - 2 \times 0.5 \times 1 \times 3$ | |
| 8(a) | $\frac{5}{7}$ oe | 1 | | |
| 8(b)(i) | $\frac{6}{35}$ oe | 2 | M1 for $\frac{6}{10}$ seen | |
| 8(b)(ii) | $\frac{33}{70}$ oe | 3 | M2 for $\frac{5}{7} \times \frac{5}{10} \times \frac{2}{7} \times \frac{4}{10}$ or M1 for one of above products | |
| 8(c) | $\frac{1250}{16807}$ | 2 | M1 for $\frac{5}{7} \times \frac{5}{7} \times \frac{5}{7} \times \frac{5}{7} \times \frac{2}{7}$ | |
| 8(d) | 4 nfww | 3 | M2 for $\frac{4}{9} \times \frac{3}{8} \times \frac{2}{7} \times \frac{5}{6}$ or M1 for a correct trial with black(s) then white | |
| 9(a)(i) | 50 | 1 | | |
| 9(a)(ii) | 24 | 2 | B1 for [l.q. =] 38 or [u.q. =] 62 seen but not as final answer | |
| 9(b) | 68 or 69 | 3 | M1 for $120 \times \frac{15}{100}$ oe (18) | |
| 10(a) | $[y=] \frac{20}{\sqrt{x}} \text{ oe}$ | 2 | M1 for reading at $(120 - their \ 18)$ seen M1 for $\frac{k}{\sqrt{x}}$ oe | |
| 10(b) | 40 | 1 | FT incorrect <i>k</i> only | |
| 10(c) | 16 | 2 | M1 <i>their</i> $20 = 5\sqrt{x}$ or better | |

| 0607/42 | Cambridge IGCSE PUBLIS | | heme October/Nove. My Mainson Partial Marks M2 for $z = K \left(\frac{their 20}{5} + 2 \right)$ | yams - |
|------------|---|-------|---|--------|
| Question | Answer | Marks | Partial Marks | OUD.C |
| 10(d) | $[z=]7\left(\frac{20}{\sqrt{x}}+2\right)$ oe final answer | 3 | M2 for $z = K\left(\frac{their20}{\sqrt{x}} + 2\right)$ or for $84 = K\left(\frac{their20}{\sqrt{4}} + 2\right)$ oe | OM |
| | | | or M1 for $z = K(y + 2)$ oe | |
| 11(a) | $-\frac{2}{5}$ oe | 2 | M1 for $5 - 7 = 2x + 3x$ oe or better | |
| 11(b) | 17 - 6x Final answer | 2 | M1 for $2(5-3x) + 7$ seen | |
| 11(c)(i) | $12 - x^2$ Final answer | 2 | M1 for $5 - 3x^2 + 2x^2 + 7$ | |
| 11(c)(ii) | $x^2 - 24x + 144$ Final answer | 3 | M1 for $(5 - 3x + 2x + 7)^2$ oe B1 for three terms of $144 - 12x - 12x + x^2$ correct | |
| 11(d) | $\frac{5-x}{3}$ oe Final answer | 2 | M1 for $x = 5 - 3y$ or $y + 3x = 5$ or $y - 5 = -3x$ or $\frac{y}{3} = \frac{5}{3} - x$ | |
| 11(e) | $\frac{13x-1}{(5-3x)(2x+7)}$ Final answer | 3 | M1 for $2(2x + 7) - 3(5 - 3x)$ or better seen M1 for common denominator (5 - 3x)(2x + 7) | |
| 12(a)(i) | Vector $\begin{pmatrix} 6\\4 \end{pmatrix}$ drawn | 1 | | |
| 12(a)(ii) | Vector $\begin{pmatrix} -2\\ 1 \end{pmatrix}$ drawn | 1 | | |
| 12(a)(iii) | Vector $\begin{pmatrix} -1\\ 4 \end{pmatrix}$ drawn | 2 | B1 for vector $\begin{pmatrix} -1\\ k \end{pmatrix}$ or vector $\begin{pmatrix} k\\ 4 \end{pmatrix}$ | |
| | | | drawn or $\begin{pmatrix} -1\\ 4 \end{pmatrix}$ seen | |

| 507/42 | Mark Scheme PUBLISHEDOctober/Nove.MarksPartial Marks $2p-q=10$ B1 | | |
|----------|---|-------|---|
| Question | Answer | Marks | Partial Marks |
| 12(b) | 2p - q = 10 3p + 4q = -7 | B1 | |
| | Equalising coefficients and adding or subtracting appropriately, or expression for one variable in terms of the other and correctly substituted, or rearranged equations seen and sketch of both lines intersecting. | M1 | |
| | p = 3, q = -4 | A2 | A1 for each If M0 A0 scored, SC1 for $p = 3$, $q = -4$ with no working or their answers satisfying one equation |