

Please check the examination details below before entering your candidate information

Candidate surname		Other names	
Pearson Edexcel International GCSE		Centre Number	Candidate Number
Wednesday 15 January 2020			
Morning (Time: 2 hours 30 minutes)		Paper Reference 4MB1/02R	
Mathematics B Paper 2R			
You must have: Ruler graduated in centimetres and millimetres, protractor, compasses, pen, HB pencil, eraser, calculator. Tracing paper may be used.			Total Marks

Instructions

- Use **black** ink or ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- **Calculators may be used.**

Information

- The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Check your answers if you have time at the end.
- Without sufficient working, correct answers may be awarded no marks.

Turn over ►

P60505A

©2020 Pearson Education Ltd.

1/1/1/1/



Answer all ELEVEN questions.

Write your answers in the spaces provided.

You must write down all the stages in your working.

1 James bought a car on 1st March 2016

On 1st March 2017, the value of the car was 20% less than the amount that James paid for the car on 1st March 2016

On 1st March 2018, the value of the car was 10% less than the value of the car on 1st March 2017

On 1st March 2018, the value of the car was £20 340

(a) Calculate the amount, in £, that James paid for the car on 1st March 2016

(3)

On 1st March 2019, the value of the car was £19 323

(b) Calculate the percentage decrease in the value of the car from 1st March 2018 to 1st March 2019

(2)

(a) In 2016, James paid £ x for his car

In 2017, the value of the car decreased by 20%.

In 2018, the value further decreased by 10% to £20 340.

So, we have to work backwards to find x .

① Find value of car in 2017, we will call this y .

$$\begin{aligned} 100\% - 10\% &= 90\% \\ 90\% &= 0.9 \end{aligned} \quad \left. \begin{array}{l} y \times 0.9 = 20\,340 \\ y = 22\,600 \end{array} \right\}$$

② Find value in 2016, x .

$$\begin{aligned} 100\% - 20\% &= 80\% \\ 80\% &= 0.8 \end{aligned} \quad \left. \begin{array}{l} x \times 0.8 = 22\,600 \\ x = £28\,250 \end{array} \right\}$$



Question 1 continued

(b)

$$\frac{\text{Percentage Change}}{\text{Change}} = \frac{\text{Original} - \text{New}}{\text{Original}} \times 100$$

$$\frac{\text{Percentage Change}}{\text{Change}} = \frac{20340 - 19323}{20340} \times 100$$
$$= 5\%$$

(Total for Question 1 is 5 marks)



- 2 A mixture for concrete is made up of 105 kg of cement, 252 kg of sand and 273 kg of gravel.

Given that

$$\text{the mass of cement} : \text{the mass of sand} : \text{the mass of gravel} = 1 : (2x + y) : (x + 2y)$$

- (a) write down two different equations in x and y .

Simplify your equations.

(2)

- (b) Hence solve these two equations for the value of x and for the value of y .

Give each value as a simplified fraction.

(3)

$$\begin{array}{ccc} \text{Cement} & : & \text{Sand} & : & \text{Gravel} \\ 105 & : & 252 & : & 273 \\ \times 105 \nearrow & & & & \nearrow \times 105 \\ 1 & : & 2x+y & : & x+2y \end{array}$$

$$105(2x+y) = 252$$

$$105(x+2y) = 273$$

$$2x+y = 2.4$$

$$x+2y = 2.6$$

- (b) The easiest way to do this will be by substitution.

$$2x+y = 2.4$$

$$y = 2.4 - 2x$$

$$x + 2(2.4 - 2x) = 2.6$$

$$x + 4.8 - 4x = 2.6$$

$$4.8 - 3x = 2.6$$

$$-3x = -\frac{11}{5}$$

$$x = \frac{11}{15}$$

$$y = 2.4 - 2\left(\frac{11}{15}\right)$$

$$y = \frac{14}{15}$$



DO NOT WRITE IN THIS AREA

Question 2 continued

Handwriting practice area with 20 horizontal dotted lines.

(Total for Question 2 is 5 marks)



- 3 (a) Find the inverse of the matrix $\begin{pmatrix} 2 & 2 \\ 5 & 4 \end{pmatrix}$

(2)

- (b) Use your answer to part (a) to find the value of x and the values of y that satisfy

$$\begin{pmatrix} 2 & 2 \\ 5 & 4 \end{pmatrix} \begin{pmatrix} y^2 - 9x \\ x \end{pmatrix} = \begin{pmatrix} 0 \\ -2 \end{pmatrix}$$

(6)

a) To find the inverse, use the formula given.

$$\begin{pmatrix} 2 & 2 \\ 5 & 4 \end{pmatrix}^{-1} = \frac{1}{(2)(4) - (2)(5)} \begin{pmatrix} (4) & -(2) \\ -(5) & (2) \end{pmatrix} = \frac{1}{-2} \begin{pmatrix} 4 & -2 \\ -5 & 2 \end{pmatrix}$$

$$= \begin{pmatrix} -2 & 1 \\ 5/2 & -1 \end{pmatrix}$$

$$\text{b) } \begin{pmatrix} y - 9x \\ x \end{pmatrix} = \begin{pmatrix} 0 \\ -2 \end{pmatrix} \div \begin{pmatrix} 2 & 2 \\ 5 & 4 \end{pmatrix} \rightarrow \div a = \times a^{-1}$$

$$\begin{pmatrix} y - 9x \\ x \end{pmatrix} = \begin{pmatrix} 0 \\ -2 \end{pmatrix} \times \begin{pmatrix} -2 & 1 \\ 5/2 & -1 \end{pmatrix}$$

To multiply a 1×2 matrix with a 2×2 matrix:

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} ax + by \\ cx + dy \end{pmatrix}$$

$$\begin{pmatrix} y - 9x \\ x \end{pmatrix} = \begin{pmatrix} 0 \\ -2 \end{pmatrix} \times \begin{pmatrix} -2 & 1 \\ 5/2 & -1 \end{pmatrix}$$

$$= \begin{pmatrix} -2(0) + 1(-2) \\ 5/2(0) + -1(-2) \end{pmatrix}$$

$$\left[\text{Inverse of matrix: } \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \right]$$



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 3 continued

$$= \begin{pmatrix} 0-2 \\ 0+2 \end{pmatrix}$$

$$= \begin{pmatrix} -2 \\ 2 \end{pmatrix}$$

(Total for Question 3 is 8 marks)



P 6 0 5 0 5 A 0 7 3 2

4 Bag B_1 and bag B_2 each contain soil.

The weight of soil in B_1 is 7.5 kg to the nearest 100 g.

The weight of soil in B_2 is 5 kg to the nearest 10 g.

(a) Find

(i) the lower bound for the total weight of soil in the two bags,

(1)

(ii) the upper bound for the difference in the weights of soil in the two bags,

(2)

(iii) the lower bound, to the nearest 50 g, for the difference in the weights of soil in the two bags.

(2)

A large jar is to be filled with sweets.

The jar will be filled using bags of sweets.

The weight of the sweets in each bag of sweets is 220 g to the nearest 5 g.

When the jar is full of sweets, the weight of sweets in the jar is 10 kg to the nearest 0.2 kg.

Bertie has 45 bags of sweets.

(b) Giving your reasons, determine whether Bertie has a sufficient number of bags of sweets to be sure that he can fill the jar with sweets.

(4)

First write bounds for each:

$$7.45 \leq B_1 < 7.55$$

$$4.995 \leq B_2 < 5.005$$

(a) (i) The lower bound is the smallest total weight of both bags:

$$7.45 + 4.995 = 12.445 \text{ kg}$$

(ii) The upper bound of the difference will be the greatest number when you subtract the two masses:

$$7.55 - 4.995 = 2.555 \text{ kg}$$

(iii) The lower bound of the difference will be the smallest number when you subtract the two masses:

$$7.45 - 5.005 = 2.445 \text{ kg} \approx 2.45 \text{ kg (nearest 50g)}$$



Question 4 continued

(b) Determine Bounds

$$217.5g \leq \text{Bag} < 222.5g$$

$$9.9kg \leq \text{Jar} < 10.1kg$$

Find maximum number of bags needed to fill jar. This is the largest number you can have when dividing weight of jar by weight of bags.

$$\frac{10.1 \times 1000}{217.5} = 46.4$$

$$46.4 > 45$$

So she does not have sufficient number of bags.



Question 4 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 4 is 9 marks)



5 Solve $\frac{3^{4x} \times 5^{3x+1} \times 45^{1-2x}}{9} = 5^4$

Show clear algebraic working.

(4)

$$\frac{3^{4x} \times 5^{3x+1} \times 45^{1-2x}}{9} = 5^4$$

$$45^{1-2x} = 5^{1-2x} \times 9^{1-2x}$$

$$\frac{3^{4x} \times 5^{3x+1} \times 5^{1-2x} \times 9^{1-2x}}{9} = 5^4$$

$$9 = 3^2$$

$$\frac{3^{4x} \times 5^{3x+1} \times 5^{1-2x} \times (3^2)^{1-2x}}{3^2} = 5^4$$

$$\text{Indices law: } (x^a)^b = x^{ab}$$

$$3^{2 \times (1-2x)} = 3^{2-4x}$$

Indices law:

$$3^{4x} \times 5^{3x+1} \times 5^{1-2x} \times 3^{2-4x} = 5^4$$

$$x^a \times x^b = x^{a+b}$$

$$3^{4x} \times 3^{2-4x} = 3^2$$

$$\frac{3^{4x+(2-4x)} \times 5^{(3x+1)+(1-2x)}}{3^2} = 5^4$$

$$5^{3x+1} \times 5^{1-2x} = 5^{x+2}$$

$$\frac{\cancel{3^2} \times 5^{x+2}}{\cancel{3^2}} = 5^4$$

$$5^{x+2} = 5^4$$

$$x^a = x^b$$

$$x+2 = 4$$

$$a = b$$

$$x = 2$$

(Total for Question 5 is 4 marks)



6 The functions f , g and h are defined as

$$f: x \mapsto x + 3$$

$$g: x \mapsto x^2 - 2x + 3$$

$$h: x \mapsto \frac{6}{x} \quad x \neq 0$$

- (a) Find (i) $g(-3)$ (ii) $fh\left(-\frac{1}{4}\right)$ (2)
- (b) (i) Express the inverse of the composite function hf in the form $(hf)^{-1}: x \mapsto \dots$ and simplify your answer. (4)
- (ii) State the value of x that must be excluded from any domain of $(hf)^{-1}$ (4)
- (c) Find the two values of x for which $hgf(x) = 2$ (5)

(a) (i) Sub $x = -3$ into $g(x)$.

$$\begin{aligned} g(-3) &= (-3)^2 - 2(-3) + 3 \\ &= 9 + 6 + 3 \\ &= 18 \end{aligned}$$

(ii) First sub $x = -\frac{1}{4}$ into $h(x)$.

$$\begin{aligned} h\left(-\frac{1}{4}\right) &= \frac{6}{\left(-\frac{1}{4}\right)} \\ &= -24 \end{aligned}$$

Next sub $x = -24$ into $f(x)$.

$$\begin{aligned} fh\left(-\frac{1}{4}\right) &= (-24) + 3 \\ &= -21 \end{aligned}$$

(b) (i) First find $hf(x)$ by substituting $x = x + 3$ into $h(x)$.

$$hf(x) = \frac{6}{x+3}$$

Next find $(hf)^{-1}: x$ by rearranging for x :

$$x(x+3) \downarrow y = \frac{6}{x+3} \uparrow \times (x+3)$$



Question 6 continued

$$\begin{aligned}
 y(x+3) &= 6 \\
 \text{Expand } \rightarrow yx + 3y &= 6 \\
 -3y \rightarrow yx &= 6 - 3y \\
 \div y \rightarrow x &= \frac{6-3y}{y} \\
 \therefore (fh)^{-1}: x &= \frac{6-3x}{x} \quad \text{Swap } y \text{ for } x
 \end{aligned}$$

(ii) Any value of the denominator being equal to 0 must be excluded.

$$\therefore x = 0$$

(c) $hgf(x) = 2$

First find $gf(x)$ by substituting $f(x)$ into $g(x)$.

$$\begin{aligned}
 gf(x) &= (x+3)^2 - 2(x+3) + 3 \\
 &= x^2 + 6x + 9 - 2x - 6 + 3 \\
 &= x^2 + 4x + 6
 \end{aligned}$$

Next find $hgf(x)$ by substituting $gf(x)$ into $h(x)$.

$$hgf(x) = \frac{6}{x^2 + 4x + 6}$$

$$\begin{aligned}
 hgf(x) &= 2 \\
 \frac{6}{x^2 + 4x + 6} &= 2 \\
 6 &= 2(x^2 + 4x + 6) \\
 6 &= 2x^2 + 8x + 12 \\
 -6 \rightarrow 0 &= 2x^2 + 8x + 6 \\
 \text{Expand } \rightarrow 0 &= 2x^2 + 8x + 6 \\
 -3 \rightarrow x+3 &= 0 \rightarrow x = -3 \\
 -1 \rightarrow x+1 &= 0 \rightarrow x = -1
 \end{aligned}$$

factorise: What 2 numbers add to make 4 and multiply to make 3? 3 and 1

(Total for Question 6 is 11 marks)



- 7 (a) Write down all the multiples of 3 lying between 1 and 20

(1)

Ahmed and Hani play a game with 20 numbered balls.
The 20 balls are numbered from 1 to 20



The balls are put in a bag.

Ahmed takes at random a ball from the bag, makes a note of its number and puts the ball back into the bag. → Always 20 in bag

If the number of the ball is a multiple of 5, Ahmed wins the game.

If Ahmed does not win the game, Hani takes at random a ball from the bag, makes a note of its number and puts the ball back into the bag.

If the number of the ball is a multiple of 3, Hani wins the game.

If Hani does not win the game, Ahmed takes at random a ball from the bag, makes a note of its number and puts the ball back into the bag.

If the number of the ball is a multiple of 6, Ahmed wins the game.

If Ahmed does not win the game, Hani takes at random a ball from the bag, makes a note of its number and puts the ball back into the bag.

If the number of the ball is a multiple of 4, Hani wins the game.

If Hani does not win the game, the game stops and the result is a draw.

The incomplete probability tree diagram for the game is shown below.


- (b) Complete the probability tree diagram for Ahmed and Hani's game.

(4)

- (c) Determine which player, Ahmed or Hani, is more likely to win the game.

Give a reason for your answer.

(5)



Question 7 continued

(a) 3, 6, 9, 12, 15, 18

(b) How many multiples of 5 between 1 and 20?

5, 10, 15, 20 $\rightarrow$ 4

$\therefore$ Probability of multiple of 5 $\rightarrow \frac{4}{20}$

How many multiples of 3 between 1 and 20?

3, 6, 9, 12, 15, 18 $\rightarrow$ 6

$\therefore$ Probability of multiple of 3 $\rightarrow \frac{6}{20}$

How many multiples of 6 between 1 and 20?

6, 12, 18 $\rightarrow$ 3

$\therefore$ Probability of multiple of 6 $\rightarrow \frac{3}{20}$

How many multiples of 4 between 1 and 20?

4, 8, 12, 16, 20 $\rightarrow$ 5

$\therefore$ Probability of multiple of 4 $\rightarrow \frac{5}{20}$

(b) Probability Ahmed wins:

$$P(A) + P(A', H', A) = \frac{4}{20} + \left(\frac{16}{20} \times \frac{14}{20} \times \frac{3}{20} \right)$$

$$= \frac{71}{250} = 0.284$$

Probability Hani wins:

$$P(A', H) + P(A', H', A', H) = \left(\frac{16}{20} \times \frac{6}{20} \right) + \left(\frac{16}{20} \times \frac{14}{20} \times \frac{17}{20} \times \frac{5}{20} \right)$$

$$= \frac{359}{1000} = 0.359$$

$$0.359 > 0.284$$

$\therefore$ Hani is more likely to win

(Total for Question 7 is 10 marks)






Figure 1 shows triangle ABC in which $AB = 100$ cm, $AC = 30$ cm and $\angle ACB = 70^\circ$

Calculate, to 3 significant figures,

- (a) the size, in degrees, of $\angle ABC$,

(3)

- (b) the length, in cm, of BC .

(3)

As shown in Figure 1, the point E lies on AB and the point D lies on AC such that ED and BC are parallel and $AD:DC = 1:2$

- (c) Write down the length, in cm, of DC .

(1)

The midpoint of BC is the point M .

- (d) Calculate the area, in cm^2 to 3 significant figures, of triangle DCM .

(2)

Given that $\frac{\text{area } \triangle ABC}{\text{area } \triangle DCM} = k$,

- (e) find the value of k without working out the area of triangle ABC .

(1)

(a) Here you should use sine rule:

$$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c}$$

$$\frac{\sin \angle ABC}{30} = \frac{\sin 70}{100}$$

$$\left[\begin{array}{l} \text{Sine rule } \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} \\ \text{Cosine rule } a^2 = b^2 + c^2 - 2bc \cos A \\ \text{Area of triangle} = \frac{1}{2} ab \sin C \end{array} \right]$$



Question 8 continued

$$\angle ABC = \sin^{-1} \left(\frac{\sin 70^\circ \times 30}{100} \right)$$

$$= 16.3741\dots$$

$$\approx \underline{\underline{16.4^\circ (3 \text{ sf})}}$$

(b) First find $\angle BAC$ and then use sine rule to find BC .

$$\angle BAC = 180 - 70 - 16.37\dots \quad (\text{Angles in a triangle add to } 180^\circ)$$
$$= 93.62\dots$$

$$\frac{BC}{\sin(93.62\dots)} = \frac{100}{\sin 70}$$

$$BC = \frac{100}{\sin 70} \times \sin(93.62\dots)$$

$$= 106.2\dots$$

$$\approx \underline{\underline{106 \text{ cm}}}$$

(c) $AD : DC \quad 30 \div (1+2) = 10$

$$\begin{array}{c} 1 : 2 \\ \swarrow \quad \searrow \\ \times 10 \quad \times 10 \\ \hline 10 : 20 \end{array}$$

$$\therefore \underline{\underline{DC = 20 \text{ cm}}}$$

(d) $\text{Area} = \frac{1}{2}ab\sin C$

$$DCM = \frac{1}{2} \times \left(\frac{1}{2} \times 106.2\dots \right) \times (20) \times \sin 70$$
$$= 498.99\dots$$

$$\approx \underline{\underline{499 \text{ cm}^2}}$$



Question 8 continued

(e) The two triangles are similar.

So $k = 3$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 8 continued

Handwriting practice area with horizontal dotted lines.

(Total for Question 8 is 10 marks)



9 The triangle A has vertices with coordinates $(-5, 2)$, $(-11, 0)$ and $(-7, 6)$.

(a) On the grid, draw and label triangle A .

(1)

Triangle B is the image of triangle A under the enlargement with centre $(1, 4)$ and scale factor $-\frac{1}{2}$.

(b) On the grid, draw and label triangle B .

(3)

Triangle C is the image of triangle B under a rotation of 180° about the point $(3, 1)$.

(c) On the grid, draw and label triangle C .

(3)

(d) Describe fully the **single** transformation that maps triangle C onto triangle A .

(3)


(b) See annotations on graph

(c) Use tracing paper to rotate B 180° about $(3, 1)$

(d) Enlargement of scale factor 2 about centre of enlargement $(9, -8)$



Question 9 continued



Turn over for a spare grid if you need to redraw your triangles.



Question 9 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA




DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 9 continued

Only use this grid if you need to redraw your triangles.



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(Total for Question 9 is 10 marks)






Figure 2

Figure 2 shows trapezium $ABCD$ in which the point O is the midpoint of AD and the point M is the midpoint of CD .

Given that $\vec{OA} = \mathbf{a}$, $\vec{CB} = \frac{1}{2}\mathbf{a}$ and $\vec{OB} = \mathbf{b}$,

(a) find, in terms of $\mathbf{a}$ and $\mathbf{b}$, simplifying your answers where possible,

- (i) $\vec{AB}$ (ii) $\vec{AC}$ (iii) $\vec{CD}$ (iv) $\vec{AM}$ (7)

The lines OB and AM intersect at N so that $\vec{AN} = \lambda \vec{AM}$ and $\vec{ON} = \mu \vec{OB}$, where λ and μ are positive constants.

(b) (i) Find and simplify an expression in terms of $\mathbf{a}$, $\mathbf{b}$ and λ for $\vec{ON}$ (2)

(ii) Hence find the value of λ and the value of μ (4)

The area of triangle OAB is 14 square units.

(c) Find the area of triangle BNA . (1)

(i) $\vec{AB} = -\vec{OA} + \vec{OB}$
 $= \mathbf{b} - \mathbf{a}$

(ii) $\vec{AC} = \vec{AB} - \vec{CB}$
 $= \mathbf{b} - \mathbf{a} - \frac{1}{2}\mathbf{a}$
 $= \mathbf{b} - \frac{3}{2}\mathbf{a}$



Question 10 continued

$$\begin{aligned} \text{(iii)} \quad \vec{CB} &= -\vec{AC} - 2(\vec{AB}) \\ &= -(b - \frac{3}{2}a) - 2a \\ &= -(\frac{1}{2}a + b) \end{aligned}$$

$$\begin{aligned} \text{(iv)} \quad \vec{AM} &= \vec{AC} + \frac{1}{2}\vec{CB} \\ &= b - \frac{3}{2}a + \frac{1}{2}(-\frac{1}{2}a - b) \\ &= -\frac{7}{4}a + \frac{1}{2}b \end{aligned}$$

$$\begin{aligned} \text{(ii)} \quad \vec{ON} &= \vec{OA} + \vec{AN} \\ &= a + \lambda \vec{AM} \\ &= a + \lambda(-\frac{7}{4}a + \frac{1}{2}b) \\ &= a + -\frac{7}{4}\lambda a + \frac{1}{2}b\lambda \\ &= a(1 - \frac{7}{4}\lambda) + \frac{\lambda}{2}b \end{aligned}$$

(iii) Since $\vec{ON}$ is a factor of $\vec{OB}$, it does not contain an a component.

$$\begin{aligned} \therefore a(1 - \frac{7}{4}\lambda) &= 0 \\ \div a \quad \rightarrow \quad 1 - \frac{7}{4}\lambda &= 0 \\ -1 \quad \rightarrow \quad -\frac{7}{4}\lambda &= -1 \\ \div -\frac{7}{4} \quad \rightarrow \quad \lambda &= \frac{4}{7} \end{aligned}$$

And the b component has to equal b .

$$\begin{aligned} \frac{\lambda}{2} &= \mu \\ \frac{\frac{4}{7}}{2} &= \mu \end{aligned}$$

$$\frac{2}{7} = \mu$$



Question 10 continued

$$(c) \text{ Area of BNA} = \frac{5}{7} \times \text{Area of OAB}$$

$$\text{Area of BNA} = \frac{5}{7} \times 14$$

$$= 10 \text{ square units}$$



DO NOT WRITE IN THIS AREA

Question 10 continued

Handwriting practice area with 20 horizontal dotted lines.

(Total for Question 10 is 14 marks)



11 Part of the curve with equation $y = x^2 - 5x + 3$ is drawn on the grid.

The equation of another curve is $y = -\frac{x^3}{6} + \frac{6x^2}{5} - \frac{3x}{2}$

(a) Complete the table of values for $y = -\frac{x^3}{6} + \frac{6x^2}{5} - \frac{3x}{2}$

Give your values of y to 2 decimal places.

x	0	0.5	1	2	3	4	4.5	5
y	0	-0.47	-0.47	0.47	1.8	2.53	2.36	1.67

(3)

(b) On the grid opposite, plot the points from your completed table and join them to form a smooth curve.

(3)

(c) Use the two curves on the grid to find an estimate, to 2 decimal places, of the range of positive values of x for which $\frac{x^3}{6} - \frac{x^2}{5} - \frac{7x}{2} + 3 < 0$

Show your working clearly.

(4)

For positive values of x , the two curves on the grid intersect at the points P and Q .

(d) Find an estimate, to 1 decimal place, of the gradient of the straight line through P and Q .

(2)

The equation of the straight line through P and Q has the form $y = ax + b$

(e) Find, to 1 decimal place, the value of b .

(2)

(a) Substitute your x -values given into the equation to find y -values.

$$y = \frac{(1)^3}{6} + \frac{6(1)^2}{5} - \frac{3(1)}{2}$$

$$y = -0.47$$

$$y = \frac{(2)^3}{6} + \frac{6(2)^2}{5} - \frac{3(2)}{2}$$

$$y = 0.47$$

$$y = \frac{(4)^3}{6} + \frac{6(4)^2}{5} - \frac{3(4)}{2}$$

$$y = 2.53$$


$$y = \frac{(4.5)^3}{6} + \frac{6(4.5)^2}{5} - \frac{3(4.5)}{2}$$

$$y = 2.36$$

(b) Plot and draw a cubic.



Question 11 continued



(c) First form the given equation using equations of two curves.

$$-\frac{x^3}{6} + \frac{6x^2}{5} - \frac{3x}{2} > x^2 - 5x + 3$$

$$-\frac{x^3}{6} + \frac{x^2}{5} + \frac{7x}{2} - 3 > 0$$

$$\frac{x^3}{6} - \frac{x^2}{5} - \frac{7x}{2} + 3 < 0$$

Now, you can find the x -values of the intersections of the two graphs.

$$0.80 < x < 4.80$$

Accepted range: $0.85 \pm 0.05 < x < 4.80 \pm 0.05$

Turn over for a spare grid if you need to redraw your curve.



Question 11 continued

$$(d) \text{ gradient} = \frac{\text{change in } y}{\text{change in } x}$$

$$\text{gradient} = \frac{1.92 - -0.5}{4.80 - 0.80} \rightarrow \text{Read values off graph}$$

$$= 0.605$$

$$\approx 0.6$$

$$(e) y = ax + b$$

$$\text{where } a = \text{gradient} = 0.605$$

$$b = y\text{-intercept}$$

$$\text{Reading from graph } b = -1.0$$




DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 11 continued

Only use this grid if you need to redraw your curve.



.....

.....

.....

.....

.....

.....

.....

.....

.....

.....



Question 11 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 11 is 14 marks)

TOTAL FOR PAPER IS 100 MARKS

