Contents Page: Number: Surds 3 7 Algebraic proofs Algebra: Transformations of graphs 13 Equations of circles 17 Quadratic and other sequences 19 Completing the square 22 Inverse and composite functions 25 Expanding more than two binomials 28 Nonlinear simultaneous equations 30 Solving quadratic inequalities 34 **Shape, Space and Measure:** Circle theorems 36 40 Vectors Sine and cosine rules 48 52 Area under graphs **Data Handling:** 57 Histograms Moving averages 65 **Probability:** Set theory 68 **Ratio and Proportion: Proportion** 71 Percentages – reverse

75

Surds

Things to remember:

- √ means square root;
- To simplify surds, find all its factors;
- To rationalise the denominator, find an equivalent fraction where the denominator is rational.

Questions:

1. Work out

$$\frac{(5+\sqrt{3})(5-\sqrt{3})}{\sqrt{22}}$$

Give your answer in its simplest form.

										(-	1	_	(0	Ì	t	ć)		,	3	3	ľ	ì	1	ĺ	2	l	ľ	•	k	(٤	>))

2. (a) Rationalise the denominator of $\frac{1}{\sqrt{3}}$

(1)

(b) Expand $(2 + \sqrt{3})(1 + \sqrt{3})$ Give your answer in the form $a + b\sqrt{3}$ where a and b are integers.

> (2) (Total 3 marks)

3. (a) Rationalise the denominator of $\frac{1}{\sqrt{7}}$

(2)

(b) (i) Expand and simplify $(\sqrt{3} + \sqrt{15})^2$ Give your answer in the form $a + b\sqrt{3}$ where a and b are integers.

.....

(ii) All measurements on the triangle are in centimetres.ABC is a right-angled triangle.k is a positive integer.




Diagram **NOT** accurately drawn

Find the value of *k*.

k =

(5)

(Total 7 marks)

Expand and simplify $(\sqrt{3} - \sqrt{2})(\sqrt{3} - \sqrt{2})$ 4. (Total 2 marks) (a) Write down the value of $49^{1/2}$ 5. (1) Write $\sqrt{45}$ in the form $k\sqrt{5}$, where k is an integer. (b) (Total 2 marks)


6. Write $\frac{\sqrt{18} + 10}{\sqrt{2}}$ in the form $a + b\sqrt{3}$ where a and b are integers.

a =

b =

(Total 2 marks)

7. Expand and simplify $(2 + \sqrt{3})(7 - \sqrt{3})$ Give your answer in the form $a + b\sqrt{3}$ where a and b are integers.



8. Rationalise the denominator of $\frac{4+\sqrt{2})(4-\sqrt{2})}{\sqrt{7}}$ Give your answer in its simplest form.

9. Show that $\frac{4-\sqrt{3})(4+\sqrt{3})}{\sqrt{13}}$ simplifies to $\sqrt{13}$

(Total for question = 2 marks)

Algebraic Proofs

Things to remember:

- Start by expanding the brackets, then factorise.
- Remember the following:
 - 1. $2n \rightarrow \text{even number}$
 - 2. $2n + 1 \rightarrow odd number$
 - 3. $a(bn + c) \rightarrow multiple of a$
 - **4.** Consecutive numbers are numbers that appear one after the other.

0		es	ti	^	n	•	
u	u	ヒコ	u	v		J	

1. (a) Expand and simplify x(x + 1)(x - 1)

(2)

In a list of three consecutive positive integers at least one of the numbers is even and one of the numbers is a multiple of 3 n is a positive integer greater than 1

(b) Prove that $n^3 - n$ is a multiple of 6 for all possible values of n.

(2)

261 – 1 is a prime number.

(c) Explain why 261 + 1 is a multiple of 3

(2) (Total for question = 6 marks)

	e that (2n + 3)² – (2n – 3)² is a multiple of 8 Il positive integer values of n.		2.
(Total for Question is 3 marks)			
	Expand and simplify (y - 2)(y - 5)	3. (a)	3.
(2)	Prove algebraically that $(2n + 1)^2 - (2n + 1)$ is an even number for all positive integer values of n.	*(b)	
(3) (Total for Question is 5 marks)			

				(3) (Total for Question is 6 marks)
		(ii)	t is a positive whole number. The expression 2t ² + 5t + 2 can never have a Explain why.	value that is a prime number.
	*(c)	(i)	Factorise 2t ² + 5t + 2	(2)
	(b)	Facto	rise y² – 10y + 16	(1)
5.	(a)	Facto	rise x² + 7x	
				(Total for Question is 4 marks)
4.	* Provintege	ve alge ers is e	braically that the difference between the squar qual to the sum of these two integers.	es of any two consecutive

6.	(a)	Factorise 3t +	12
0.	(a)	i actorise st	1 4

(1)

(b) (i) Expand and simplify
$$7(2x + 1) + 6(x + 3)$$

.....


(ii) Show that when x is a whole number 7(2x + 1) + 6(x + 3) is always a multiple of 5

(3) (Total for Question is 4 marks)

7. Prove that
$$(n-1)^2 + n^2 + (n+1)^2 = 3n^2 + 2$$

(Total for Question is 2 marks)

8.	Prove algebraically that the difference between the squares of any two consecutive integers is equal to the sum of these two integers.
	(Total for question is 4 marks)
9.	The product of two consecutive positive integers is added to the larger of the two integers. Prove that the result is always a square number.
	(Total for question = 3 marks)
	11



Transformations of graphs


Things to remember:

- 1. f(x) means the function of x.
- 2. -f(x) is a reflection in the x-axis.
- 3. f(-x) is a reflection in the y-axis.
- 4. f(x a) is a translation in the x-axis, a units.
- 5. f(x) + b is a translation in the y-axis, b units.
- 6. cf(x) is an enlargement in the y-axis, scale factor c.
- 7. f(dx) is an enlargement in the x-axis, scale factor $\frac{1}{d}$.

Questions:


1. y = f(x)

The graph of y = f(x) is shown on the grid.



(a) On the grid above, sketch the graph of y = -f(x).

The graph of y = f(x) is shown on the grid.



The graph **G** is a translation of the graph of y = f(x).


(b) Write down the equation of graph **G**.

......


(Total for Question is 3 marks)

(2)

2. The graph of y = f(x) is shown on both grids below.




(a) On the grid above, sketch the graph of y = f(-x)




(b) On this grid, sketch the graph of y = -f(x) + 3

(1) (Total for question = 2 marks)

- The graph of y = f(x) is shown on each of the grids. (a) On this grid, sketch the graph of y = f(x 3)3.




On this grid, sketch the graph of y = 2f(x)(b)



(Total for Question is 4 marks)


(2)

The graph of y = f(x) is shown on the grid. 4.



On the grid above, sketch the graph of y = f(x + 3)(a)

(2) The graph of y = g(x) is shown below.



The graph **G** is the reflection of y = g(x) in the *x*-axis. (b) Write down an equation of graph **G**.

(1)

(Total for question = 3 marks)


Equations of Circles

Things to remember:

- 8. The general equation of a circle is $(x a)^2 + (y b)^2 = r^2$, where (a, b) is the centre and r is the radius.
- 9. To calculate the equation of the tangent:
 - 1. Calculate the gradient of the radius of the circle.
 - 2. Calculate the gradient of the tangent of the circle.
 - 3. Substitute the given coordinate and the gradient of the tangent into y = mx + c to calculate the y-intercept.

Questions:

- 1. The circle C has radius 5 and touches the y-axis at the point (0, 9), as shown in the diagram.
 - (a) Write down an equation for the circle C, that is shown in the diagram.



(3)

A line through the point P(8, -7) is a tangent to the circle C at the point T.

(b) Find the length of PT.

(3) (Total 6 marks)

2.	A circle C has centre (-1, 7) and passes through the p	point (0, 0). Find an equation for C.
		(Total 4 marks)
3.	The circle C has centre (3, 1) and passes through the	point P(8, 3).
	(a) Find an equation for C.	
		(4)
	(b) Find an equation for the tangent to C at P.	
		(5) (Total 9 marks)

Quadratic and Other Sequences

Th	ina	s to	rem	Δm	hΔ	r.
111	my	ร เบ	16111	IGIII	ne	Ι.

_	s to remember: To calculate the nth term of a quadratic sequence: 1. Calculate the first difference. 2. Calculate the second difference. 3. How many n²s? 4. Subtract. 5. Calculate the nth term of the difference. 6. Write the quadratic nth term.
Quest 1.	Here are the first 5 terms of a quadratic sequence. 1 3 7 13 21 Find an expression, in terms of <i>n</i> , for the <i>n</i> th term of this quadratic sequence.
	(Total for question is 3 marks)
2.	Here are the first six terms of a Fibonacci sequence. 1 1 2 3 5 8 The rule to continue a Fibonacci sequence is, the next term in the sequence is the sum of the two previous terms. (a) Find the 9th term of this sequence.

The first three terms of a different Fibonacci sequence are a + b

Show that the 6th term of this sequence is 3a + 5b(b)

(2)

(1)

	n that the 3rd term is 7 and the 6th term is 29, find the value of <i>a</i> and the value of <i>b</i> .	
(3) (Total for question = 6 marks	b =	
	are the first five terms of a sequence. 2 8 18 32 50 Find the next term of this sequence.	
(1	 nth term of a different sequence is $3n^2 - 10$ Work out the 5th term of this sequence.	The <i>n</i> t (b)
(1 (Total for question = 2 marks		

h term of this sequence.	Here are the first five terms of an arithmetic sequence. 1 5 9 13 17 (a) Write down an expression, in terms of <i>n</i> , for the <i>n</i> th	4.
(2)	The n th term of a different number sequence is $3n^2 + 7$ (b) Find the 10th term of this sequence.	
(2) (Total for Question is 4 marks)		

Completing the Square

Things to remember:

- To complete the square:
 - 1. Halve the coefficient of x.
 - 2. Put in brackets with the x and square the brackets.
 - 3. Subtract the half-coefficient squared.
 - 4. Don't forget the constant on the end!
 - 5. Simply.
- For $(x p)^2 + q = 0$, the turning point is (p, q).

Questions:


1. (i) Sketch the graph of $f(x) = x^2 - 5x + 10$, showing the coordinates of the turning point and the coordinates of any intercepts with the coordinate axes.

(ii) Hence, or otherwise, determine whether f(x + 2) - 3 = 0 has any real roots. Give reasons for your answer.

(Total for question = 6 marks)

2.	(a)	Write $2x^2 + 16x + 35$ in the form $a(x + b)^2 + c$ where a , b , and c are integers.
	(b)	Hence, or otherwise, write down the coordinates of the turning point of the graph of $y = 2x^2 + 16x + 35$
		(1)
3.	The e (a)	(Total for question = 4 marks) xpression $x^2 - 8x + 21$ can be written in the form $(x - a)^2 + b$ for all values of x . Find the value of a and the value of b .
		a =
		b =(3)

The equation of a curve is y = f(x) where $f(x) = x^2 - 8x + 21$ The diagram shows part of a sketch of the graph of y = f(x).



The minimum point of the curve is *M*. (b) Write down the coordinates of *M*.

(Total for Question is 4 marks)

Inverse and Composite Functions

Things to remember:

- y = f(x) means that y is a function of x.
- f(a) means the value of x is a, so substitute x with a.
- The graph of the inverse is the reflection of the graph in the line y = x
- We find the inverse function by putting the original function equal to y and rearranging to make x the subject.
- We use the notation f-1(x) for the inverse function.
- When a function is followed by another, the result is a composite function.
- fg(x) means do g first, followed by f.

Questions:

1. The functions f and g are such that

$$f(x) = 1 - 5x$$
 and $g(x) = 1 + 5x$

(a) Show that gf(1) = -19

(b) Prove that
$$f^{-1}(x) + g^{-1}(x) = 0$$
 for all values of x .

(2)

2.	The f	Function f is such that $f(x) = 4x - 1$ Find $f^{-1}(x)$	
		function g is such that g(x) = kx² where k is a constant. In that fg(2) = 12 work out the value of k	$f^{-1}(x) = \dots$ (2)
3.	The f	functions f and g are such that $f(x) = 3(x - 4)$ and $g(x) = \frac{x}{5} + 1$ Find the value of $f(10)$	k =(2) (Total for question = 4 marks)
	(b)	Find $g^{-1}(x)$	(1)
			$g^{-1}(x) = \dots$ (2)

Show that ff(x) = 9x - 48(c)

(Total for question = 5 marks)

 $f(x) = 3x^2 - 2x - 8$ Express f(x + 2) in the form $ax^2 + bx$ 4.

(Total for question is 3 marks)

Expanding more than two binomials

Things to remember:

- Start by expanding two pair of brackets using the grid or FOIL method.
- Then expand the third set of brackets.
- Use columns to keep x³, x² etc in line to help with addition.

_				4 =				
11	ш	Δ	e	tı	\sim	n	e	•
Q	u	ᆫ	J	u	v		J	

1. Show that

$$(x-1)(x+2)(x-4) = x^3 - 3x^2 - 6x + 8$$

for all values of x.

(Total for question is 3 marks)

2. Show that

$$(3x - 1)(x + 5)(4x - 3) = 12x^3 + 47x^2 - 62x + 15$$

for all values of x.

(Total for question is 3 marks)

3.	Show that	9
	$(x-3)(2x+1)(x+3) = 2x^3 + x^2 - 18x - 18x$	
	for all values of x	f

(Tot	al for qu	uestion	is 3 r	marks)

4. $(2x + 1)(x + 6)(x - 4) = 2x^3 + ax^2 + bx - 24$ for all values of x, where a and b are integers. Calculate the values of a and b.

Nonlinear Simultaneous Equations

Things to remember:

- 1. Substitute the linear equation into the nonlinear equation.
- 2. Rearrange so it equals 0.
- 3. Factorise and solve for the first variable (remember there will be two solutions).
- 4. Substitute the first solutions to solve for the second variable.
- 5. Express the solution as a pair of coordinate where the graphs intersect.

Questions:

1.	Solve the equations
	$x^2 + y^2 = 36$
	x = 2y + 6

(Total for Question is 5 marks)

3. Solve the simultaneous equations $x^2 + y^2 = 25$ y = 2x + 5

$x = \dots$	and $y = $
	0
$x = \dots \dots$	and $y = $
	(Total for Question is 6 marks

4. Solve the simultaneous equations $x^2 + y^2 = 9$ x + y = 2Give your answers correct to 2 decimal places.

	x =											<i>y</i> :	=												
or .	x =											<i>y</i> :	=												
				(To	ot	a	۱ :	f	01	• (Qι	ıe	S	ti	0	n	į	S	6	r	n	aı	'n	S

5.	Solve algebraically the simultaneous equations $x^2 + y^2 = 25$ y - 2x = 5
	y-2x=5
	(Total for Question is 5 marks)
	22

Solving Quadratic Inequalities

Things to remember:

• Start by solving the quadratic to find the values of x, then sketch the graph to determine the inequality.

Questions:

1. Solve $x^2 > 3x + 4$

(Total for question = 3 marks)

2. Solve the inequality $x^2 > 3(x+6)$

(Total for question = 4 marks)


3. Solve the inequality $x^2 + 5x > 6$

(Total for question = 3 marks)


4.	Solve the inequality	$x^2 - 2x + 8 < 0$	
			(Total for question = 3 marks)
_			, , , , , , , , , , , , , , , , , , , ,
5.	Solve the inequality	x² - x ≥ 12	
			(Total for question = 3 marks)
6.	Solve the inequality	$x^2 \le 4(2x + 5)$	
•	corre and anoquality	x = 1(= x = 0)	
			(Total for question = 4 marks)

Circle theorems


Things to remember:




The angle at the centre is twice the angle at the circumference.




The angle in a semicircle is 90°.




Angles subtended by the same arc are equal.




Opposite angles in a cyclic quadrilateral sum to 180°.



Tangents from a point are equal.



A tangent is perpendicular to a radius.



Angles in alternate segments are equal.

Questions:

1.




Diagram **NOT** accurately drawn

P is a point on the circumference of the circle, centre *O*.

PQ is a tangent to the circle.

(i) Write down the size of angle *OPQ*.

(ii) (Give a reason	for your	answer.
--------	---------------	----------	---------

(Total 2 marks)




Diagram **NOT** accurately drawn

A, B and C are points on the circumference of a circle, centre O.

AC is a diameter of the circle.

(a)	(i)	Write down the size of angle <i>ABC</i> .

(ii)	Give a reason for your answer.




Diagram **NOT** accurately drawn

D, E and F are points on the circumference of a circle, centre O. Angle $DOF = 130^{\circ}$.

- (b) (i) Work out the size of angle *DEF*.
 - (ii) Give a reason for your answer.

(2) (Total 4 marks)

(2)

3.




Diagram **NOT** accurately drawn

A and B are points on the circumference of a circle, centre O.

PA and PB are tangents to the circle.

Angle APB is 86°.

Work out the size of the angle marked x.

° (Total 2 marks)

4.




Diagram **NOT** accurately drawn

In the diagram, A, B, C and D are points on the circumference of a circle, centre O.

Angle $\overrightarrow{BAD} = 70^{\circ}$. Angle $\overrightarrow{BOD} = x^{\circ}$.

Aligic BOB - X .

Angle $BCD = y^{\circ}$.


(a)	(1)	work out the value of x.	

(11)	Give a reason for your answer.

(b) (i) Work out the value of *y*.

(ii)	Give a reason for your answer.	

(2) (Total 4 marks) 5.



DA is a tangent to the circle. Angle $ADO = 36^{\circ}$

(a)	Work out the size of angle AOD.	
-----	---------------------------------	--

(2)	Work out the size of angle <i>ABC</i> .	(i)	(b)
	Give a reason for your answer.	(ii)	
(3) (Total 5 marks)			

Vectors

Things to remember:

- Use the letter provided in the question.
- Going against the arrow is a negative.
- Vectors need to be written in bold or underlined.
- They can be manipulated similarly to algebra.

Questions:

1. The diagram shows a regular hexagon *ABCDEF* with centre *O*.




Diagram **NOT** accurately drawn

 $\overrightarrow{OA} = 6\mathbf{a}$ $\overrightarrow{OB} = 6\mathbf{b}$

(a) Express in terms of **a** and/or **b**

(i) \overrightarrow{AB} ,

(ii) \overrightarrow{EF} .

.....

.....

(2)

X is the midpoint of BC.


(b) Express \overrightarrow{EX} in terms of **a** and/or **b**

(2)

Y is the point on AB extended, such that AB : BY = 3:2

(c) Prove that E, X and Y lie on the same straight line.

(3) (Total 7 marks) 2. T is the point on PQ for which PT : TQ = 2 : 1.



OPQ is a triangle.

$$\overrightarrow{OP}$$
 = **a** and \overrightarrow{OQ} = **b**.

(a) Write down, in terms of **a** and **b**, an expression for \overrightarrow{PQ} .

\overrightarrow{PQ}	=	 	 		 _	 				 		_		 			_		 _	 						
		 	 	 -	 -		-	•		-		-			•	 	-	-		 •	 	-	,	('	1	1

(b) Express \overrightarrow{OT} in terms of **a** and **b**. Give your answer in its simplest form.

$$\overrightarrow{OT} = \dots$$
 (2) (Total 3 marks)

3. *OABC* is a parallelogram.




Diagram **NOT** accurately drawn

P is the point on *AC* such that $AP = \frac{2}{3}AC$.

$$\overrightarrow{OA} = 6a. \overrightarrow{OC} = 6c.$$

(a) Find the vector \overrightarrow{OP} . Give your answer in terms of **a** and **c**.

																								((3	

The midpoint of CB is M.

(b) Prove that *OPM* is a straight line.

(2) (Total 5 marks) OPQ is a triangle.R is the midpoint of OP.S is the midpoint of PQ.

$$\overrightarrow{OP}$$
 = **p** and \overrightarrow{OQ} = **q**




Diagram **NOT** accurately drawn

(i) Find \overrightarrow{OS} in terms of **p** and **q**.

 \overrightarrow{OS} =

(ii) Show that RS is parallel to OQ.

(Total 5 marks)

OPQR is a trapezium with PQ parallel to OR. 5.

$$\overrightarrow{OP}$$
 = 2**b**

$$\overrightarrow{PQ}$$
 = 2a \overrightarrow{OR} = 6a

$$OR = 6a$$

M is the midpoint of PQ and N is the midpoint of OR.




Diagram NOT accurately drawn

Find the vector \overrightarrow{MN} in terms of **a** and **b**. (a)

$$\overrightarrow{MN} =$$
 (2)

X is the midpoint of MN and Y is the midpoint of QR.

Prove that XY is parallel to OR. (b)

(Total 4 marks)

6. ABCD is a straight line.




Diagram **NOT** accurately drawn

O is a point so that $\overrightarrow{OA} = \mathbf{a}$ and $\overrightarrow{OB} = \mathbf{b}$.

B is the midpoint of AC.

C is the midpoint of AD.


Express, in terms of \mathbf{a} and \mathbf{b} , the vectors

 \overrightarrow{AC} (i)

(ii)	OD
(11)	

(Total 3 marks)

7. Diagram **NOT** accurately drawn



ABCD is a parallelogram. AB is parallel to DC.

AD is parallel to BC.

$$\overrightarrow{AB} = \mathbf{p}$$


$$\stackrel{
ightarrow}{AD}$$
 = q

Express, in terms of p and q (a)

- \overrightarrow{AC} (i)
- \overrightarrow{BD} (ii)

(2)

Diagram **NOT** accurately drawn



AC and BD are diagonals of parallelogram ABCD. AC and BD intersect at T.

Express \overrightarrow{AT} in terms of **p** and **q**. (b)

(Total 3 marks)


8.	Diagram	NOT	accurately	drawr
	_		_	

OAB is a triangle.

B is the midpoint of *OR*.

Q is the midpoint of AB.

$$\overrightarrow{OP}$$
 = 2a \overrightarrow{PA} = a \overrightarrow{OB} = b



- (a) Find, in terms of **a** and **b**, the vectors
 - (i) \overrightarrow{AB} ,
 - (ii) \overrightarrow{PR} ,
 - (iii) \overrightarrow{PQ}

	(III) ~ ~ .	
′h\	Hence explain why POP is a straight line	(4)

(b) Hence explain why *PQR* is a straight line.

(2)

The length of PQ is 3 cm.

(c) Find the length of *PR*.

.....cm (1)

(Total 7 marks)


Sine and Cosine Rules

Things to remember:

• For any triangle ABC, $a^2 = b^2 + c^2 - 2bc \cos A$

For any triangle ABC, $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$

For any triangle ABC, area = ½ a b sinC



Questions:

1. Diagram **NOT** accurately drawn

ABC is a triangle.

D is a point on AC.

Angle $BAD = 45^{\circ}$

Angle $ADB = 80^{\circ}$


AB = 7.4 cm

DC = 5.8 cm

Work out the length of BC.

Give your answer correct to 3

significant figures.



(Total for question = 5 marks)

2. Diagram **NOT** accurately drawn

ABC is a triangle.

AB = 8.7 cm.


Angle $ABC = 49^{\circ}$.

Angle $ACB = 64^{\circ}$.

Calculate the area of triangle ABC.

Give your answer correct to 3 significant

figures.



(Total for Question is 5 marks)

ABCD is a quadrilateral.
 Diagram NOT accurately drawn
 Work out the length of DC.
 Give your answer correct to 3 significant figures.






Diagram NOT accurately drawn
 ABC is an isosceles triangle.
 Work out the area of the triangle.
 Give your answer correct to 3 significant figures.




.....cm² (Total for Question is 4 marks)

5. Diagram NOT accurately drawn The diagram shows triangle LMN. Calculate the length of LN. Give your answer correct to 3 significant figures.



(Total for Question is 5 marks)


6. VABCD is a solid pyramid. ABCD is a square of side 20 cm. The angle between any sloping edge and the plane ABCD is 55° Calculate the surface area of the pyramid. Give your answer correct to 2 significant figures.



.....cm²
(Total for question = 5 marks)

7. The diagram shows triangle ABC. The area of triangle ABC is $k\sqrt{3}$ cm².

Find the exact value of *k*.




k	=			
		(Total for o	question =	7 marks)

8. Diagram **NOT** accurately drawn

AC = 9.2 mBC = 14.6 m

Angle ACB = 64°

(a) Calculate the area of the triangle ABC.Give your answer correct to 3 significant figures.



(b) Calculate the length of *AB*. Give your answer correct to 3 significant figures.

(3)

(2)

(Total for Question is 5 marks)

Area under Graphs

Things to remember:

- Velocity is speed with direction
- Acceleration and deceleration is given by the gradient of the graph $\left(\frac{rise}{run}\right)$
- The distance travelled is given by the area under the graph.

Questions:

1. A car has an initial speed of u m/s.

The car accelerates to a speed of 2u m/s in 12 seconds.


The car then travels at a constant speed of 2u m/s for 10 seconds.

Assuming that the acceleration is constant, show that the total distance, in metres, travelled by the car is 38u.

(Total for question = 4 marks)

2. Karol runs in a race.


The graph shows her speed, in metres per second, *t* seconds after the start of the race.



(a) Calculate an estimate for the gradient of the graph when t = 4 You must show how you get your answer.

(3)	Describe fully what your answer to part (a) represents.	(b)
(2)		, ,
	Explain why your answer to part (a) is only an estimate.	(c)
(1) otal for question = 6 marks)	(Т	

3. Here is a speed-time graph for a car journey. The journey took 100 seconds.




The car travelled 1.75km in the 100 seconds.

(a) Work out the value of V.

(3)	Describe the acceleration of the car for each part of this journey.	(b)
(2) uestion = 5 marks)	(Total for qu	

4. The graph shows information about the velocity, *v* m/s, of a parachutist *t* seconds after leaving a plane.




(a) Work out an estimate for the acceleration of the parachutist at t = 6

 m/s²
(2)

(b) Work out an estimate for the distance fallen by the parachutist in the first 12 seconds after leaving the plane.
Use 3 strips of equal width.

m
(3
(Total for question is 5 marks

5. Here is a speed-time graph for a car.



(a) Work out an estimate for the distance the car travelled in the first 10 seconds. Use 5 strips of equal width.

Is your answer to (a) an underestimate or an overestimate of the actual distance? Give a reason for your answer.
(1)
(Total for question = 4 marks)

Histograms

Things to remember:

- Frequency = Frequency Density x Class Width;
- The y-axis will always be labelled "frequency density";
- The x-axis will have a continuous scale.


Questions:

1. One Monday, Victoria measured the time, in seconds, that individual birds spent on her bird table. She used this information to complete the frequency table.

Time (t seconds)	Frequency
0 < <i>t</i> ≤ 10	8
10 < <i>t</i> ≤ 20	16
20 < t ≤ 25	15
25 < <i>t</i> ≤ 30	12
30 < <i>t</i> ≤ 50	6


(a) Use the table to complete the histogram.





(3)

On Tuesday she conducted a similar survey and drew the following histogram from her results.




Time (Seconds)

(b) Use the histogram for Tuesday to complete the table.

Time (t seconds)	Frequency
0 < <i>t</i> ≤ 10	10
10 < <i>t</i> ≤ 20	
20 < <i>t</i> ≤ 25	
25 < <i>t</i> ≤ 30	
30 < <i>t</i> ≤ 50	

(2) (Total 5 marks) 2. This histogram gives information about the books sold in a bookshop one Saturday.



(a) Use the histogram to complete the table.


Price (P) in pounds (£)	Frequency
$0 < P \le 5$	
5 < <i>P</i> ≤ 10	
10 < <i>P</i> ≤ 20	
20 < <i>P</i> ≤ 40	

(2)

The frequency table below gives information about the books sold in a second bookshop on the same Saturday.

Price (P) in pounds (£)	Frequency
0 < <i>P</i> ≤ 5	80
5 < <i>P</i> ≤ 10	20
10 < <i>P</i> ≤ 20	24
20 < <i>P</i> ≤ 40	96


(b) On the grid below, draw a histogram to represent the information about the books sold in the second bookshop.



(3)

(Total 5 marks)

3. The incomplete table and histogram give some information about the distances walked by some students in a school in one year.



Use the information in the histogram to complete the frequency table. (a)

Distance (d) in km	Frequency
0 < <i>d</i> ≤ 300	210
300 < <i>d</i> ≤ 400	350
400 < <i>d</i> ≤ 500	
500 < <i>d</i> ≤ 1000	

density

(2)

Use the information in the table to complete the histogram. (b)


(Total 3 marks)

4. The incomplete histogram and table show information about the weights of some containers.

Weight (w) in kg	Frequenc y
$0 < w \le 1000$	16
1000 < <i>w</i> ≤ 2000	
2000 < <i>w</i> ≤ 4000	
4000 < <i>w</i> ≤ 6000	16
6000 < <i>w</i> ≤ 8000	
8000 < <i>w</i> ≤ 12000	8

(a) Use the information in the histogram to complete the table.


(b) Use the information in the table to complete the histogram.



(Total 4 marks)

(2)

5. The incomplete histogram and table give some information about the distances some teachers travel to school.



(a) Use the information in the histogram to complete the frequency table.

Distance (dkm)	Frequency
$0 < d \le 5$	15
5 < <i>d</i> ≤ 10	20
10 < <i>d</i> ≤ 20	
20 < <i>d</i> ≤ 40	
40 < <i>d</i> ≤ 60	10

(2)


(b) Use the information in the table to complete the histogram.

(Total 3 marks)


6. The table gives information about the heights, in centimetres, of some 15 year old students.

Height (h cm)	145 < <i>h</i> ≤ 155	155 < <i>h</i> ≤ 175	175 < <i>h</i> ≤ 190
Frequency	10	80	24

Use the table to draw a histogram.



7. A teacher asked some year 10 students how long they spent doing homework each night. The histogram was drawn from this information.



Use the histogram to complete the table.

Time (<i>t</i> minutes)	Frequency
10 ≤ <i>t</i> < 15	10
15 ≤ <i>t</i> < 30	
30 ≤ <i>t</i> < 40	
40 ≤ <i>t</i> < 50	
50 ≤ <i>t</i> < 70	

(Total 2 marks)

Moving Averages

Things to remember:

- In this context, averages means the mean (add the numbers and divide by how many there were.
- Moving averages are used to identify trends in data peaks, troughs, increasing and decreasing trends.

Questions:

1. The table shows the number of computer games sold in a supermarket each month from January to June.

Jan	Feb	Mar	Apr	May	Jun
147	161	238	135	167	250

Work out the three month moving averages for this information.

2. The table shows the number of digital cameras Bytes sold each month in the first six months of 2005.

Month	January	February	March	April	May	June
Number of digital	30	19	20	15	27	39
cameras sold						

The first 3-month moving average for this data is 23

Work out the **second** 3-month moving average for this data.

(Total 2 marks)

3. The table shows the number of orders received each month by a small company.

Month	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug
Number of								
orders	23	31	15	11	19	16	20	13
received								

Work out the first two 4-month moving averages for this data.

an	d
(Total 3 marks)

4. A shop sells DVD players.

The table shows the number of DVD players sold in every three-month period

The table shows the number of DVD players sold in every three-month period from January 2003 to June 2004.

Year	Months	Number of DVD players sold
2003	Jan – Mar	58
	Apr – Jun	64
	Jul – Sep	86
	Oct – Dec	104
2004	Jan – Mar	65
	Apr – Jun	70

	Apr = Juri	
(a)	Calculate the set of four-point moving averages for this data.	
(b)	What do your moving averages in part (a) tell you about the trend	(2) in the sale of DVD
	players?	
		(1)
Paul a	and Carol open a new shop in the High Street.	(1 (Total 3 marks
The to	able shows the monthly takings in each of the first four months	

MonthJanFebMarchAprilMonthly takings (£)9375890792559420

5.

Work out the 3-point moving averages for this information.


 (Total 2 marks)

1110	Year		lonths	om Januai Num	ber of CE]		
	2004		an – Mar	270	DCI OI OL				
	2001		pr – Jun	324					
			ul – Sept	300					
			ct – Dec	258					
	2005		an – Mar	309					
		Α	pr – Jun	335			=		
(a)	Calc	ulate the d	complete	set of four	-point mo	ving aver	ages for th	is info	ormation.
(b)	Whai	t trend do	these mo	oving aver	ages sugg	est?			
(b)	What	t trend do	these mo	oving avera	ages sugç	est?			
(b)	What	t trend do	these mo	oving avera	ages sugg	est?			
(b)	What	t trend do	these mo	oving avera	ages sugg	est?			
(b)	What	t trend do	these mo	oving avera	ages sugg	est?			(Total 3 r
				oving avera					(Total 3 r
	 table sh		ne informa		t student a		Summer 2005		(Total 3 r

434, 443,	,		
	(Total	2 marks)	

Set Theory

Things to remember:




The **intersection** is where two sets overlap.

$$A \cap B$$

This means **A** and **B**.






If you put two sets together, you get the union.

$$A \cup B$$

This means **A** or **B**.





The **complement of A** is the region that is not A.

A'

This means **not A**.

Questions:

1.

$$\mathcal{E}$$
= {1, 2, 3, 4, 5, 6, 7, 8, 9, 10}

$$A = \{\text{multiples of 2}\}\$$

$$A \cap B = \{2, 6\}$$

$$A \cup B = \{1, 2, 3, 4, 6, 8, 9, 10\}$$

Draw a Venn diagram for this information.


(Total for question is 4 marks)

- 2. Here is a Venn diagram.
 - Write down the numbers (a) that are in set
 - (i) $A \cup B$

.....

 $A \cap B$ (ii)

..... (2)



One of the numbers in the diagram is chosen at random.


Find the probability that the number is in set A'

(Total for question = 4 marks)

- 3. Here is a Venn diagram.
 - Write down the numbers that (a) are in set
 - $A \cup B$ (i)

(ii) $A \cap B$

..... (2)



One of the numbers in the diagram is chosen at random.

(b) Find the probability that the number is in set A'

(Total for question = 4 marks)

Sami asked 50 people which drinks they liked from tea, coffee and milk. All 50 people like at least one of the drinks 19 people like all three drinks. 16 people like tea and coffee but do not like milk. 21 people like coffee and milk. 24 people like tea and milk. 40 people like coffee. 1 person likes only milk. Sami selects at random one of the 50 people. (a) Work out the probability that this person likes tea.
(b) Given that the person selected at random from the 50 people likes tea, find the probability that this person also likes exactly one other drink.
(2) (Total for question = 6 marks)

4.

Proportion

Things to remember:

- Start by checking the question for squares, cubes and roots;
- "x is directly proportional to y" looks like x α y or x = ky
- "x is inversely proportional to y" looks like $\mathbf{x} \propto \frac{1}{y}$ or $\mathbf{x} = \frac{k}{y}$

\sim			4:	_		_	_
u	u	es	iti	O	n	S	Ξ

- 1. The shutter speed, S, of a camera varies inversely as the square of the aperture setting, f. When f = 8, S = 125
 - (a) Find a formula for S in terms of f.

(3)

(b) Hence, or otherwise, calculate the value of S when f = 4

S =(1)

(Total 4 marks)

2. In a factory, chemical reactions are carried out in spherical containers. The time, *T* minutes, the chemical reaction takes is directly proportional to the square of the radius, *R* cm, of the spherical container.

When R = 120, T = 32

Find the value of T when R = 150

T =(Total 4 marks)

	directly proportional to the square of <i>t</i> . 80 when <i>t</i> = 4	
	Express <i>d</i> in terms of <i>t</i> .	(a)
(3)	Work out the value of <i>d</i> when <i>t</i> = 7	(b)
(1)	$d = \dots$ Work out the positive value of t when $d = 45$	(c)
(2) (Total 6 marks)		4 . The 0
ional to the square of the time, t,	e distance, <i>D</i> , travelled by a particle is directly proportion en. When <i>t</i> = 40, <i>D</i> = 30 Find a formula for <i>D</i> in terms of <i>t</i> .	
(3)	$D = \dots$ Calculate the value of D when $t = 64$	(b)
(1)	Calculate the value of <i>t</i> when <i>D</i> = 12 Give your answer correct to 3 significant figures.	(c)
(2) (Total 6 marks)		


	to the mass of water, m kg, in the water heater. W (a) Find T when $m = 400$	5.
	The time, T seconds, it takes a water heater to be proportional to the power, P watts, of the water he When $P = 1400$, $T = 360$ (b) Find the value of T when $P = 900$	
T =(3) (Total 6 marks) conds.	A ball falls vertically after being dropped. The ball falls a distance <i>d</i> metres in a time of <i>t</i> se <i>d</i> is directly proportional to the square of <i>t</i> . The ball falls 20 metres in a time of 2 seconds. (a) Find a formula for <i>d</i> in terms of <i>t</i> .	6.
d =(3)	c) Calculate the distance the ball falls in 3 seconds.	(b)
m (1)	Calculate the time the ball takes to fall 605 m.	(c)
seconds (3) (Total 7 marks)		

7.		pring, the tension (T newtons) is directly proportional to its extension (x cm). When the on is 150 newtons, the extension is 6 cm. Find a formula for T in terms of x .
		<i>T</i> =
	(b)	Calculate the tension, in newtons, when the extension is 15 cm.
		newtons (1)
	(c)	Calculate the extension, in cm, when the tension is 600 newtons.
		cm
0	fic in	(1) (Total 5 marks)
8.	When	versely proportional to d . a $d = 50$, $f = 256$ he value of f when $d = 80$
		f =(Total 3 marks)

Percentages - reverse

Things to remember:

• Work out what the multiplier would have been;



Ques 1.	After	nsulation reduces annual heating costs by 20 he insulated his loft, Curtley's annual heating cout Curtley's annual heating cost would have	g cost was £520.
			£(Total 3 marks)
2.	Andre The s	sale, normal prices are reduced by 20%. SALE - 20% OFF ew bought a saddle for his horse in the sale. sale price of the saddle was £220. ulate the normal price of the saddle.	
			£(Total 3 marks)
3.	This Bill s	a's weekly pay this year is £240 is 20% more than her weekly pay last year. ays 'This means Hajra's weekly pay last year s wrong, Explain why.	was £192'.
	(b)	Work out Hajra's weekly pay last year.	(1)

£

(Total 3 marks)

4.	(a) The price of all rail season tickets to London increase (a) The price of a rail season ticket from Cambrid Work out the price before this increase.	
	(b) After the increase, the price of a rail season to £2828.80 Work out the price before this increase.	£(2) icket from Brighton to London was
		£
		(3) (Total 5 marks)
5.	In a sale, normal prices are reduced by 25%. The sale price of a saw is £12.75 Calculate the normal price of the saw.	
		£
6.	In a sale, normal prices are reduced by 12%. The sale price of a DVD player is £242. Work out the normal price of the DVD player.	(Total 3 marks)
		£
7.	A garage sells cars. It offers a discount of 20% off the normal price for carbave pays £5200 cash for a car. Calculate the normal price of the car.	(Total 3 marks)
		£
		(Total 3 marks)