

**Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries**

Q	Working	Answer	Mark	Notes
<b>1</b> (b)	$(-5)^2 - 4 \times -5$ oe e.g. $25 + 20$		2	M1 for a correct substitution
		45		A1
<b>2</b> (c)	$5x - 3 = 4(2x + 3)$ oe <b>or</b> $\frac{5x}{4} - \frac{3}{4} = 2x + 3$ oe		3	M1 for correctly removing the denominator, condone missing brackets
	e.g. $5x - 8x = 12 + 3$ <b>or</b> $-3x = 12 + 3$ <b>or</b> $8x - 5x = -12 - 3$ <b>or</b> $3x = -12 - 3$ <b>or</b> $-\frac{3}{4} - 3 = 2x - \frac{5x}{4}$ <b>or</b> $-\frac{15}{4} = \frac{3x}{4}$			M1 for a correct rearrangement with terms in $x$ on one side and numbers on the other, allow correct rearrangement of their equation in the form $ax + b = cx + d$
		-5		A1 dep on at least M1  SCB2 for an answer of $x = -2$ coming from $5x - 3 = 8x + 3$ <b>or</b> $x = 5$ coming from $5x - 3 = 2x + 12$
				<b>Total 3 marks</b>
<b>3</b> (a)		0	1	B1 condone $150^0$
	(b)	-2	1	B1 condone $3^{-2}$
				<b>Total 2 marks</b>

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
4	e.g. $35x + 10y = 27.5$ <b>or</b> $21x + 6y = 16.5$ $\frac{6x - 10y = 34}{41x} = 61.5$ $\frac{21x - 35y = 119}{41y} = -102.5$  e.g. $3x - 5\left(\frac{5.5 - 7x}{2}\right) = 17$ or $7\left(\frac{17 + 5y}{3}\right) + 2y = 5.5$ oe		4	M1 for a correct method to eliminate $x$ or $y$ :  coefficients of $x$ or $y$ the same <b>and</b> correct operator to eliminate selected variable (condone any one arithmetic error in multiplication)  <b>or</b> writing $x$ or $y$ in terms of the other variable and correctly substituting.
		$x = 1.5$ <b>or</b> $y = -2.5$		A1 oe, dep on M1
				M1 (dep on 1 <sup>st</sup> M1) for a correct method to find other variable by substitution of found variable into one equation  <b>or</b> for repeating the above method to find the second variable.
		$x = 1.5$ <b>and</b> $y = -2.5$		A1 oe, dep on M1
				<b>Total 4 marks</b>

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
5 (a)			2	M1 for $(x \pm 6)(x \pm 7)$
		$(x + 6)(x - 7)$		A1 for $(x + 6)(x - 7)$ or $(x - 7)(x + 6)$ isw roots given if candidate solves the quadratic = 0
(b)	$3x - 8x < 3 - 15$ or $15 - 3 < 8x - 3x$		3	M1 accept as equation or with the wrong inequality sign.
	$-5x < -12$ or $12 < 5x$			M1 accept as equation or with the wrong inequality sign.
		$x > 2.4$		A1 Accept $2.4 < x$ or $x > \frac{12}{5}$ oe allow $(-\infty, 2.4)$  award M1 M1 A0 for 2.4 with = sign or no inequality or incorrect inequality sign.
				<b>Total 5 marks</b>

**Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries**

Q	Working	Answer	Mark	Notes
6	$(4x+1)(x-3) = 4x^2 - 12x + x - 3 (= 4x^2 - 11x - 3)$ $(4x+1)(5x+6) = 20x^2 + 24x + 5x + 6 (= 20x^2 + 29x + 6)$ $(x-3)(5x+6) = 5x^2 + 6x - 15x - 18 (= 5x^2 - 9x - 18)$		3	M1 for multiplying 2 brackets with at least 3 out of 4 terms correct
	$(5x+6)(4x^2 - 11x - 3) = 20x^3 - 55x^2 - 15x + 24x^2 - 66x - 18$ $(x-3)(20x^2 + 29x + 6) = 20x^3 + 29x^2 + 6x - 60x^2 - 87x - 18$ $(4x+1)(5x^2 - 9x - 18) = 20x^3 - 36x^2 - 72x + 5x^2 - 9x - 18$			M1 (dep) for multiplying the product of the first 2 brackets (ft from the 1 <sup>st</sup> stage) by the 3 <sup>rd</sup> bracket, and getting at least 3 out of 6 or 4 out of 8 terms correct
		$20x^3 - 31x^2 - 81x - 18$		A1
	<b>Alternative</b>			
	$20x^3 + 24x^2 - 60x^2 + 5x^2 - 15x + 6x - 72x - 18$			B2 for at least 6 out of 8 terms correct (B1 for 4 or 5 out of 8 correct terms)
		$20x^3 - 31x^2 - 81x - 18$		A1
				<b>Total 3 marks</b>

**Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries**

Q	Working	Answer	Mark	Notes
7 (a)		9, 28, 45, 63, 76, 80	1	B1
(b)			2	B2 for a correct cf graph with points at ends of intervals and joined with a curve or line segments If not B2 then B1 for 5 or 6 of their points (ft from a table with only one arithmetic error) at ends of intervals and joined with a curve or line segments <b>OR</b> for 5 or 6 points plotted correctly at ends of intervals not joined <b>OR</b> for 5 or 6 of their points from table plotted consistently within each interval (not at upper ends of intervals) at their correct heights and joined with a curve or line segments
(c)	e.g. reading across from 40 and reading down		2	M1 ft reading from a cf graph provided method is shown
		35 - 38		A1 ft from their cf graph
				<b>Total 5 marks</b>

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
8	e.g. $\frac{15}{4}$		3	M1 for $3\frac{3}{4}$ expressed as an improper fraction
	e.g. $\frac{15^5}{4} \times \frac{7}{9^3}$ OR $\frac{105}{36}$ oe			M1 correct cancelling or multiplication of numerators and denominators without cancelling
	e.g. $\frac{15^5}{4} \times \frac{7}{9^3} = \frac{35}{12} = 2\frac{11}{12}$ or $\frac{15}{4} \times \frac{7}{9} = \frac{105}{36} = \frac{35}{12} = 2\frac{11}{12}$ or $\frac{15}{4} \times \frac{7}{9} = \frac{105}{36} = 2\frac{33}{36} = 2\frac{11}{12}$	shown		A1 dep on M2, for conclusion to $2\frac{11}{12}$ from correct working – either sight of the result of the multiplication e.g. $\frac{105}{36}$ oe must be seen or correct cancelling prior to the multiplication to $\frac{35}{12}$ NB: use of decimals scores no marks
				<b>Total 3 marks</b>
9 (a)		33.75	1	B1 oe eg 33.750
(b)		33.85	1	B1 allow 33.849̇ or 33.849 <sup>r</sup> or “33.8499...” do NOT allow 33.879 without indication of recurring “9”
				<b>Total 2 marks</b>

**Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries**

Q	Working	Answer	Mark	Notes
<b>10</b> (a) (i)		40	2	B1 cao (may be written on the diagram)
(ii)		<u>Angles in same segment</u> (are equal)		B1 <b>or</b> <u>angles at the circumference from the same arc</u> of the circle <b>or</b> <u>angles on the same arc</u> of the circle <b>Alternatively:</b> (two applications of) <u>Opposite angles of a cyclic quadrilateral</u> sum to 180°
(b)		140	1	B1 cao (may be written on the diagram)
				<b>Total 3 marks</b>

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
11 (a)		(-4.5) 3 4.5 (3) 1.5 (3) 10.5	2	B2 for all correct (B1 for any two correct) No points in table but correctly plotted on grid, award mark
(b)	(-3, -4.5) (-2,3) (-1,4.5) (0,3) (1,1.5) (2,3) (3,10.5)	Smooth curve	2	B2 for a correct smooth curve. Points or curve passing through correct values within half a small square. (B1 for at least 5 points plotted correctly; ft from table for plotting only provided B1 awarded in part (a))
(c)			2	M1 for drawing $y = -x - 1$ with two correct points plotted and intersection with curve. <b>or</b> for stating $y = -x - 1$ <b>or</b> for $\frac{1}{2}x^3 - 2x + 3 = -x - 1$ seen
		-2.3 to -2.4		A1 ft their curve dep on M1 <b>and</b> line $y = -x - 1$ drawn
				<b>Total 6 marks</b>



Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
12	$yn^2 = n^2 + d$ or $y = 1 + \frac{d}{n^2}$		4	M1
	$yn^2 - n^2 = d$ or $-d = n^2 - yn^2$ or $y - 1 = \frac{d}{n^2}$			M1
	$n^2(y - 1) = d$ or $-d = (1 - y)n^2$			M1 for factorising $n^2$ from a suitable expression. or $n^2 = \frac{d}{y - 1}$
		$n = \sqrt{\frac{d}{y - 1}}$		A1 Accept $n = \sqrt{\frac{-d}{1 - y}}$ Penalise $\pm\sqrt{\quad}$
				<b>Total 4 marks</b>

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
13 (a)	$T = \frac{k}{m^2}$ or $Tm^2 = k$		3	M1 for a correct equation with a constant Do not allow constant = 1
	$30 \times 0.5^2 = k$ or $30 = \frac{k}{0.5^2}$ or $k = 7.5$ or $k = \frac{15}{2}$			M1 dep on M1 for correct substitution in a correct equation M2 for $k = 7.5$ or $k = \frac{15}{2}$
		$T = \frac{7.5}{m^2}$		A1 for $T = \frac{7.5}{m^2}$ or $T = \frac{15}{2m^2}$ SCB2 for $Tm^2 = 7.5$ or $Tm^2 = \frac{15}{2}$ or $m^2 = \frac{7.5}{T}$ or $m^2 = \frac{15}{2T}$
(b)		750	1	B1 cao
				<b>Total 4 marks</b>

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
14			2	M1 Arcs on $BC$ , $AB$ and arcs from these points meeting <b>or</b> for bisector without arcs
		Correct bisector		A1 must see correct arcs
				<b>Total 2 marks</b>

15	$-3(x^2 - 4x) + 7$ or $-3\left(x^2 - 4x - \frac{7}{3}\right)$		4	M1 for factorising the expression to find $b$ or $b = -3$ stated or shown clearly in answer.
	$-3[(x-2)^2 \dots]$ or $c = -2$ $-3[(x-2)^2 - 4] + 7$ or $-3\left[(x-2)^2 - 4 - \frac{7}{3}\right]$			M1 or for $c$ shown clearly in answer.
	$-3(x-2)^2 + 12 + 7$ or $-3\left[(x-2)^2 - \frac{19}{3}\right]$			M1 fully correct method.
		$19 - 3(x-2)^2$		A1 for $19 - 3(x-2)^2$ oe
				<b>Total 4 marks</b>

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
16 (a)	$(f(2.6) =) 5 \times 2.6 - 7 (= 6)$ <b>or</b> $gf(x) = \frac{5(5x-7)}{5x-7+4}$ oe		2	M1 for finding $f(2.6)$ <b>or</b> $gf(x)$
		3		A1
(b)	$5\left(\frac{5x}{x+4}\right) - 7 = 2$ <b>or</b> $\frac{5x}{x+4} = \frac{2+7}{5}$ oe		3	M1
	$25x = 9(x+4)$ oe			M1 for removing the denominator $(x+4)$ in a correct equation
		2.25		A1 oe
ALT (b)	$fg(x) = 2 \Rightarrow g(x) = f^{-1}(2) (=9/5)$ and attempt at $f^{-1}$ or $f^{-1}(2)$			M1
	$x = g^{-1}(“9/5”)$			M1
		2.25		A1 oe
(c)	$y = \frac{5x}{x+4}$ <b>or</b> $x = \frac{5y}{y+4}$ $y(x+4) = 5x$ $x(y+4) = 5y$		3	M1
	e.g. $4y = x(5-y)$ <b>or</b> e.g. $4x = y(5-x)$			M1 for a correct rearrangement and factorising
		$\frac{4x}{5-x}$		A1 oe e.g. $\frac{-4x}{x-5}$
				<b>Total 8 marks</b>

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
17	$(y-4)^2 - (y-4) + y^2 = 10$ or $x^2 - x + (x+4)^2 = 10$		6	M1 for substituting linear equation into the quadratic equation
	$2y^2 - 9y + 10 = 0$ or $2x^2 + 7x + 6 = 0$			A1 for a correct equation in the form $ax^2 + bx + c = 0$ or $ax^2 + bx = -c$ or equations of the same form but in $y$
	$(2y-5)(y-2) = 0$ or $\frac{-9 \pm \sqrt{(-9)^2 - (4 \times 2 \times 10)}}{2 \times 2}$ or $(2x+3)(x+2) = 0$ or $\frac{-7 \pm \sqrt{7^2 - (4 \times 2 \times 6)}}{2 \times 2}$			M1ft For solving <i>their</i> 3 term quadratic equation using any correct method. If factorising, allow brackets which expanded give 2 out of 3 terms correct (if using formula or completing the square allow one sign error and some simplification – allow as far as eg $\frac{-7 \pm \sqrt{49 - 48}}{4}$ or eg $\left(x + \frac{7}{4}\right)^2 - \frac{1}{16} = 0$ oe $\frac{9 \pm \sqrt{81 - 80}}{4}$ or eg $\left(y - \frac{9}{4}\right)^2 - \frac{1}{16} = 0$ oe
	$(-1.5, 2.5)$ and $(-2, 2)$			A1 for both pairs of coordinates oe eg $\left(\frac{-3}{2}, \frac{5}{2}\right)$ accept coordinates listed as pairs, ie $x_1, y_1, x_2, y_2$
	$\sqrt{('1.5' - '2')^2 + ('2.5' - '2')^2}$			M1 dep on M1 for finding length of AB
		$\frac{\sqrt{2}}{2}$		A1 dep M3
				<b>Total 6 marks</b>

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
18		$BDF = 70^\circ$	4	B1 may be marked on diagram
		<u>Alternate segment</u> theorem		B1 reason, the angle between a tangent and a chord is equal to the angle subtended in the <u>alternate segment</u>
		$EFB = 180 - (70 + 40) = 70$ <u>opposite angles</u> in a <u>cyclic quadrilateral</u>		B1 Angle $EFB$ with reason, <u>opposite angles</u> in a <u>cyclic quadrilateral</u> sum to $180^\circ$
		$CBF = EFB$ <u>alternate</u> angles therefore $EF$ is parallel to $ABC$		B1 conclusion, <u>alternate</u> angles are equal
				<b>Total 4 marks</b>

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
19	$AB = 2b - 2a$ or $BA = 2a - 2b$ $MN = 10a - b$ or $NM = -10a + b$		5	M1 for finding $AB$ or $BA$ or $MN$ or $NM$
	eg $MP = -b + 2a + k(2b - 2a)$ and $MP = l(10a - b)$ or eg $MP = b + k(2a - 2b)$ and $MP = l(10a - b)$ or eg $PN = 8a + k(2a - 2b)$ and $PN = l(10a - b)$ or eg $AP = 8a + k(b - 10a)$ and $AP = l(2b - 2a)$ or eg $AP = -2a + b + k(10a - b)$ and $AP = l(2b - 2a)$ or eg $AM = k(2b - 2a) + l(b - 10a)$ and $AM = -2a + b$			M2 for writing eg $MP$ or $PN$ or $AP$ or $AM$ in two different ways in terms of $a$ and $b$ (M1 for writing eg $MP$ or $PN$ or $AP$ or $AM$ in one way)  These may be written as eg $PM$ in place of $MP$
	eg $2 - 2k = 10\lambda$ and $-1 + k = -\lambda$ (from $MP$ 1st) or eg $2k = 10\lambda$ and $1 - 2k = -\lambda$ (from $MP$ 2nd) or eg $8 + 2k = 10\lambda$ and $-2k = -\lambda$ (from $PN$ ) or eg $8 - 10k = -2\lambda$ and $k = 2\lambda$ (from $AP$ 1st) or eg $-2 + 10k = -2\lambda$ and $1 - k = 2\lambda$ (from $AP$ 2nd) or eg $-2k - 10\lambda = -2$ and $2k + \lambda = 1$ (from $AM$ )			M1 dep M3 for writing a pair of equations using their variables. $MP$ (1st) leads to $\lambda = \frac{1}{9}, k = \frac{4}{9}$ $MP$ (2nd) leads to $\lambda = \frac{1}{9}, k = \frac{5}{9}$ $PN$ leads to $\lambda = \frac{8}{9}, k = \frac{4}{9}$ $AP$ (1st) leads to $\lambda = \frac{4}{9}, k = \frac{8}{9}$ $AP$ (2nd) leads to $\lambda = \frac{4}{9}, k = \frac{1}{9}$ $AM$ leads to $\lambda = \frac{1}{9}, k = \frac{4}{9}$
		4 : 5		A1 cao

Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
20	$12^2 = 2^4 \times 3^2$ or $2 \times 12^2 = 2^5 \times 3^2$ oe or $\frac{2 \times 12^2}{3^2} (= 32) = 2^5$		5	M1
	$18^{4n} = (2 \times 3^2)^{4n}$ or $2^{4n} \times 3^{2 \times 4n}$			M1
	$3n^2 - 14n - 5 (= 0)$			A1
	e.g. $(3n + 1)(n - 5) (= 0)$  $n = \frac{14 \pm \sqrt{(-14)^2 - (4 \times 3 \times -5)}}{2 \times 3}$			M1 for solving their 3 term quadratic equation using any correct method - if factorising, allow brackets which expanded give 2 out of 3 terms correct (if using formula or completing the square allow one sign error and some simplification – allow as far as e.g. $\frac{14 \pm \sqrt{196 + 60}}{6}$ oe)
		$-\frac{1}{3}, 5$		A1 Allow $-0.33$ or better for $-\frac{1}{3}$
				<b>Total 5 marks</b>



## Practice Tests Set 15 – Paper 1H mark scheme, performance data and suggested grade boundaries

Q	Working	Answer	Mark	Notes
---	---------	--------	------	-------

Qn	Skill tested	Mean score	Max score	Mean %	Edexcel averages: scores of candidates who achieved grade:							
					ALL	9	8	7	6	5	4	3
1	Expressions and formulae	1.69	2	85	1.69	1.99	1.94	1.80	1.64	1.27	0.57	0.00
2	Linear equations	2.70	3	90	2.70	2.98	2.88	2.75	2.74	2.35	2.11	0.33
3	Powers and roots	1.67	2	84	1.67	1.96	1.79	1.73	1.49	1.38	1.03	0.33
4	Simultaneous linear equations	3.31	4	83	3.31	3.95	3.85	3.42	2.89	2.38	1.39	0.67
5	Inequalities	4.04	5	81	4.04	4.89	4.59	4.07	3.77	2.61	1.96	1.11
6	Algebraic manipulation	2.35	3	78	2.35	2.96	2.68	2.38	2.04	1.69	0.64	0.11
7	Statistical measures	3.89	5	78	3.89	4.75	4.43	3.95	3.68	2.68	1.53	0.22
8	Fractions	2.21	3	74	2.21	2.64	2.30	2.36	2.17	1.62	1.11	0.33
9	Degree of accuracy	1.31	2	66	1.31	1.89	1.55	1.32	0.79	0.58	0.18	0.00
10	Angles, lines and triangles	1.93	3	64	1.93	2.46	2.21	1.98	1.45	1.16	0.93	0.11
11	Graphs	3.78	6	63	3.78	4.78	3.87	3.57	3.17	2.95	2.25	1.11
12	Algebraic manipulation	2.47	4	62	2.47	3.78	3.17	2.15	1.40	0.55	0.25	0.11
13	Proportion	2.22	4	56	2.22	3.49	2.60	1.79	1.24	0.83	0.36	0.00
14	Construction	1.09	2	55	1.09	1.68	1.33	0.89	0.55	0.50	0.11	0.00
15	Quadratic equations	1.95	4	49	1.95	3.51	2.15	1.50	0.64	0.18	0.18	0.00
16	Function notation	4.31	8	54	4.31	7.42	5.18	1.40	1.17	0.32	0.44	0.00
17	Quadratic equations	2.49	6	42	2.49	5.11	2.61	1.08	0.53	0.14	0.11	0.00
18	Geometrical reasoning	0.97	4	24	0.97	1.97	1.02	0.48	0.09	0.13	0.07	0.00
19	Vectors	0.67	5	13	0.67	1.37	0.74	0.35	0.04	0.04	0.00	0.00
20	Powers and roots	1.37	5	27	1.37	3.28	1.02	0.31	0.06	0.06	0.00	0.00
	<b>TOTAL</b>	<b>46.42</b>	<b>80</b>	<b>58</b>	<b>46.42</b>	<b>66.86</b>	<b>51.91</b>	<b>39.28</b>	<b>31.55</b>	<b>23.42</b>	<b>15.22</b>	<b>4.43</b>

### Suggested grade boundaries

Grade	9	8	7	6	5	4	3
Mark	60	46	36	28	19	10	2