



There is a nice way that you can remember the difference between the volume and surface area of a sphere if you have covered integration and differentiation. The integral of the surface area of a sphere is equal to the volume and the derivative of the volume gives you the surface area.


 $\int \text{ surface area of a sphere} = \int 4\pi r^2 dr = \frac{4\pi r^3}{3} = \text{volume of a sphere} \qquad \text{and} \qquad \text{derivative of volume of a sphere} = \frac{d}{dr} \left(\frac{4\pi r^3}{3}\right) = 4\pi r^2 = \text{surface area of a sphere}$ Going off topic from 3D shapes, did you know that you can do the same thing for 2D shapes with the area of a circle to get the circumference?  $A = \pi r^2 \Rightarrow \frac{dA}{dr} = 2\pi r$ 


Another cool fact: The volume of a cone + volume of sphere = volume of a cylinder (if all 3 shapes have the same height and diameter of course)


## **Examples**















Area of cross section = area of blue shaded semi-circle + area of green shaded rectangle =  $\frac{1}{2}\pi(30)^2 + 60(45) = 4113.7 \ cm^2$ Volume = area of cross section ×depth =  $4113.7176 \times 90 = 370,234.5 \ cm^3$ 






and have used straight away instead





Find the volume of the frustrum



Since we are given the dimensions of both cones we can use the fact that it is volume of the bigger cone minus the volume of the smaller cone, so all you need to know is the formula for the volume of a cone  $\frac{1}{3}\pi r^2 h$ 


Here it looks like we don't have enough info as the radius of the cone is missing. We use similar shapes to get r. These shapes are similar so


$$\frac{40}{15} = \frac{20}{r}$$


$$40r = 300$$

$$r = 7.5$$
Volume  $= \frac{1}{3}\pi (15)^2 (40) - \frac{1}{3}\pi (7.5)^2 (20) = 5890.2 \ cm^3$ 

## What happens when we have two shapes together such as cones, spheres or cylinders?








Total volume =  $136.7 \ cm^3$ 


When finding surface area remember that we care about the areas of everything on the OUTSIDE. The white bottom circle of the cylinder is not exposed and neither is white circle of the cone. Surface area of cylinder=  $2\pi(3)(3.5) + \pi(3)^2 = 30\pi$ 

Surface area of cone=  $\pi(3)(5) = 15\pi$ (note: it is not  $+\pi r^2$  since the circle is not exposed) Total surface area = 141.4  $m^2$ 

## Working backwards:

Sometimes we are given the volume/surface area of the of the shape. We can use this to work backwards and solve for r (or another unknown). Once we have r we can find the volume of the hemisphere and then add the volume together to find the total volume.





$$\pi r(4) + \pi r^{2} = \frac{33}{4}\pi$$

$$4\pi r + \pi r^{2} = \frac{33}{4}\pi$$

$$4r + r^{2} = \frac{33}{4}\pi$$


$$4r + r^{2} = \frac{33}{4}\pi$$


$$4r + r^{2} = \frac{33}{4}\pi$$

$$4r^{2} + 16r - 33 = 0$$

$$(2r + 11)(2r - 3) = 0$$


$$r \neq -\frac{11}{2}, r = \frac{3}{2}$$







## Algebraic Side Lengths:


The diagram below shows a cylinder and a sphere. The radius of the base of the cylinder is 2x cm and the height of the cylinder is h cm. The radius of the sphere 3x cm. The volume of the cylinder is equal to the volume of the sphere. Express h in terms of x Volume of cylinder =  $\pi r^2 h = \pi (2x)^2 (h) = \pi (4x) (h) = 4\pi h x^2$ Volume of sphere=  $\frac{4}{3}\pi r^3 = \frac{4}{3}\pi (27x^3)$ Volumes are equal  $\Rightarrow 4\pi h x^2 = \frac{4}{3}\pi (27x^3)$ Cancel the  $\pi$  on both sides  $4hx^2 = \frac{4}{3}(27x^3)$   $4hx^2 = 36x^3$ h = 9x








$$\frac{1}{3}\pi r^2 h = \frac{1}{3}\pi (15)^2 (36) = 2700\pi$$
  
k = 2700







 $\pi(12x)(20x) = 2160\pi$   $240x^{2} = 2160$   $x^{2} = 9$ x = 3

Now we know x we can find the volume of the cone

Volume of cone  $\frac{1}{3}\pi r^2 h = \frac{1}{3}\pi (12x)^2 (16x) = \frac{1}{3}\pi (12\times 3)^2 (16\times 3) = 20736\pi$ 





 $\frac{1}{3}x^{2}(4x) + \frac{2}{3}x^{3} = (2x)^{2}h$  $\frac{4}{3}x^{2}(x) + \frac{2}{3}x^{3} = 4x^{2}h$  $4x^{2}(x) + 2x^{3} = 3(4x^{2}h)$  $6x^{3} = 12x^{2}h$  $12x^{2}h = 6x^{3}$  $h = \frac{6x^{3}}{12x^{2}}$  $h = \frac{x}{2}$